

Research

Open Access

Effect of time of administration on cholesterol-lowering by psyllium: a randomized cross-over study in normocholesterolemic or slightly hypercholesterolemic subjects

Guido MA Van Rosendaal*^{1,2}, Eldon A Shaffer², Alun L Edwards² and Rollin Brant¹

Address: ¹Department of Community Health Sciences, University of Calgary, 3330 Hospital Drive NW, Calgary, Alberta, Canada, T2N 4N1 and ²Department of Medicine, University of Calgary, 3330 Hospital Drive NW, Calgary, Alberta, Canada, T2N 4N1

Email: Guido MA Van Rosendaal* - gvrosend@ucalgary.ca; Eldon A Shaffer - shaffer@ucalgary.ca; Alun L Edwards - aedwards@ucalgary.ca; Rollin Brant - brant@ucalgary.ca

* Corresponding author

Published: 28 September 2004

Received: 05 May 2004

Nutrition Journal 2004, **3**:17 doi:10.1186/1475-2891-3-17

Accepted: 28 September 2004

This article is available from: <http://www.nutritionj.com/content/3/1/17>

© 2004 Van Rosendaal et al; licensee BioMed Central Ltd.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Reports of the use of psyllium, largely in hypercholesterolemic men, have suggested that it lowers serum cholesterol as a result of the binding of bile acids in the intestinal lumen. Widespread advertisements have claimed an association between the use of soluble fibre from psyllium seed husk and a reduced risk of coronary heart disease. Given the purported mechanism of cholesterol-lowering by psyllium, we hypothesized that there would be a greater effect when psyllium is taken with breakfast than when taken at bedtime. Secondly, we expected to confirm a cholesterol-lowering effect of psyllium in subjects with "average" cholesterol levels.

Methods: Sixteen men and 47 women ranging in age from 18 to 77 years [mean 53 +/- 13] with LDL cholesterol levels that were normal or slightly elevated but acceptable for subjects at low risk of coronary artery disease were recruited from general gastroenterology and low risk lipid clinics. Following a one month dietary stabilization period, they received an average daily dose of 12.7 g of psyllium hydrophilic mucilloid, in randomized order, for 8 weeks in the morning and 8 weeks in the evening. Change from baseline was determined for serum total cholesterol, LDL, HDL and triglycerides.

Results: Total cholesterol for the "AM first" group at baseline, 8 and 16 weeks was 5.76, 5.77 and 5.80 mmol/L and for the "PM first" group the corresponding values were 5.47, 5.61 and 5.57 mmol/L. No effect on any lipid parameter was demonstrated for the group as a whole or in any sub-group analysis.

Conclusion: The timing of psyllium administration had no effect on cholesterol-lowering and, in fact, no cholesterol-lowering was observed. Conclusions regarding the effectiveness of psyllium for the prevention of heart disease in the population at large may be premature.

Background

A cholesterol lowering effect has been reported for a variety of soluble dietary fibres [1-5]. In February, 1998, the U.S. Food and Drug Administration authorized the use, on food labels and food labelling, of health claims on the association between soluble fibre from psyllium seed husk and a reduced risk of coronary heart disease [6]. Among the suggested mechanisms by which soluble fibre lowers cholesterol is the binding of bile acids in the intestinal lumen resulting in decreased absorption and increased faecal excretion of them [7-15]. The ensuing bile acid depletion increases hepatic demand for the de novo synthesis of bile acids from cholesterol. This requirement is met, in part, by increased hepatic LDL receptor activity, which in turn reduces circulating LDL.

The accumulation and concentration of bile in the gallbladder is a continuous process. In rats which lack a gallbladder, the biliary excretion rate of bile salts is maximal at night [16] and bile is stored in the gallbladder during an overnight fast. To the extent that the cholesterol-lowering effect of psyllium requires an interaction with bile, the magnitude of its cholesterol-lowering effect should vary with the quantity of bile in contact with a given amount of psyllium. The gallbladder empties with a meal and the total quantity of bile salt presented to the small bowel should be highest when this emptying occurs at breakfast, following a fast and the associated overnight accumulation of secreted bile. Conversely, psyllium presented to the gut following a short fast and without the full stimulus to gallbladder emptying of an associated meal should result in an encounter with a smaller amount of bile. We accordingly set out to test the hypothesis that administering psyllium with breakfast would have a significantly greater cholesterol-lowering effect than would taking a similar dose at bedtime. Since the predominance of the literature with regard to the cholesterol-lowering effect of psyllium is in individuals with hypercholesterolemia, a secondary goal of our study was to confirm the cholesterol-lowering effect of psyllium in subjects with "average" cholesterol levels.

Methods

Patients identified in gastroenterology practices as requiring long-term treatment with psyllium, typically for chronic constipation or the irritable bowel syndrome, were invited to participate in the study. In addition, individuals who had received dietary counselling in a lipid clinic regarding cholesterol-lowering and who subsequently had cholesterol levels deemed not to require further intervention because they met targets set out in clinical practice guidelines were invited by clinic staff to participate. Subjects were deemed ineligible if they were under age 18 years, were under active treatment for hyperlipidemia, had total cholesterol greater than 7.00 mmol/

L, required alterations in dosage of medications which might have an effect on lipid levels, had had a gastrectomy, had any disease which is associated with hyperlipidemia, were receiving a bile acid binding resin, or if they did not eat breakfast regularly.

The study was approved by the Conjoint Health Research Ethics Board of the University of Calgary, Faculty of Medicine. Subjects were given a description of the study indicating our interest in comparing the relative efficacy of hs versus am dosing with psyllium without indicating the specific hypothesis, and were asked to sign a consent form.

Gastroenterology patients were given a high fibre diet sheet as part of their therapeutic regimen and were asked to take this on a continuing basis, beginning one month prior to the initiation of the study. The diet sheet emphasized dietary sources containing predominantly insoluble fibre. Lipid clinic patients had all received in-depth counselling regarding dietary measures for hypercholesterolemia, including a high fibre regimen, and had implemented their dietary changes at least one month before beginning the study. An unsweetened psyllium preparation, "Novo-Mucilax" [NovoPharm], providing three grams of hydrophilic mucilloid per 6.2 gram powder, and a scoop known to provide at least ten grams of psyllium were provided. Containers were numbered and weighed at the conclusion of each test interval. Subjects were randomized to initially take a scoop full of psyllium either with breakfast or at bedtime. Using a crossover design, psyllium was taken in the morning or evening for eight weeks and at the alternate time for the subsequent eight weeks.

Determinations of serum total cholesterol, LDL, HDL and triglycerides were made before beginning psyllium, at eight weeks and at sixteen weeks after commencing its use. A trained dietician obtained a dietary history and patients were weighed at the beginning and at the conclusion of the study. All lipid determinations were undertaken following a 14 hour fast and analyses were done in a central laboratory.

After initial data inspection based on boxplots and summary measures, cholesterol values at 8 and 16 weeks were examined using analysis of variance, taking into account treatment, period, and between and within subject effects in accordance with the cross-over design. The pattern of change from baseline to 16 weeks was evaluated using paired t-tests.

Results

Of 86 subjects beginning the study, 33 of those referred from the gastroenterology clinics and 30 of those referred

from the lipid clinic completed it. Of those withdrawing, eight did so because they could not tolerate the psyllium or it was felt to interfere with prescribed medication, 2 had elevated lipids, 4 did not complete all the required blood work, 5 were unable to comply with the protocol because of work or lifestyle changes and 4 developed intercurrent diseases which precluded completing the protocol.

The age range of subjects was 18 to 77 years, mean 53 +/- 13 years, including 16 men and 47 women. The mean

dose of psyllium taken was 12.7 +/- 2.3 g for morning dosing and 12.7 +/- 2.2 g when taken in the evenings.

Values for total, LDL, HDL cholesterol and triglycerides at baseline, eight and sixteen weeks for various subgroupings are tabulated in tables 1, 2, 3 and 4, respectively. Data for changes from baseline for total, LDL and HDL cholesterol and triglycerides are listed in table 5. Changes from baseline were not significant for any parameter. Furthermore, the confidence intervals given in Table 5 show that this data essentially excludes a clinically significant psyllium effect.

Table 1: Total Cholesterol in mmol/L [SD]

[n]	Baseline	8 wks	16 wks
AM first	5.76 [0.94]	5.77 [1.01]	5.80 [0.95]
PM first	5.47 [0.98]	5.61 [1.10]	5.57 [1.06]
Females [47]	5.72 [0.96]	5.79 [1.07]	5.74 [0.99]
Males [16]	5.29 [0.94]	5.39 [0.96]	5.48 [1.04]
GI subjects [33]	5.00 [0.83]	5.06 [0.91]	5.07 [0.84]
Lipid clinic Subjects [30]	6.28 [0.59]	6.39 [0.69]	6.35 [0.70]
All subjects [63]	5.61 [0.96]	5.69 [1.05]	5.68 [1.00]

"AM first" are subjects taking psyllium in the morning for the first 8 weeks and in the evening for the second 8 weeks. Vice versa for "PM first".

Table 2: LDL CHOLESTEROL in mmol/L [SD]

	Baseline	8 wks	16 wks
AM first	3.55 [1.01]	3.51 [1.04]	3.52 [0.94]
PM first	3.39 [0.90]	3.43 [1.03]	3.45 [0.92]
Females	3.55 [0.92]	3.54 [1.04]	3.52 [0.92]
Males	3.22 [1.03]	3.24 [0.99]	3.39 [0.97]
GI subjects	2.89 [0.81]	2.85 [0.85]	2.91 [0.72]
Lipid clinic subjects	4.09 [0.66]	4.15 [0.72]	4.11 [0.69]
All subjects	3.47 [0.95]	3.47 [1.03]	3.48 [0.92]

Table 3: HDL CHOLESTEROL in mmol/L [SD]

	Baseline	8 wks	16 wks
AM first	1.38 [0.29]	1.42 [0.37]	1.42 [0.33]
PM first	1.38 [0.40]	1.39 [0.36]	1.36 [0.33]
Females	1.41 [0.30]	1.44 [0.30]	1.42 [0.29]
Males	1.28 [0.45]	1.29 [0.50]	1.30 [0.43]
GI subjects	1.36 [0.39]	1.41 [0.43]	1.37 [0.38]
Lipid clinic subjects	1.40 [0.30]	1.40 [0.28]	1.41 [0.27]
All subjects	1.38 [0.35]	1.41 [0.36]	1.39 [0.33]

Table 4: TRIGLYCERIDES in mmol/L [SD]

	Baseline	8 wks	16 wks
AM first	1.81 [0.70]	1.84 [0.86]	1.86 [0.91]
PM first	1.53 [0.52]	1.69 [0.70]	1.65 [0.61]
Females	1.66 [0.61]	1.72 [0.70]	1.76 [0.71]
Males	1.70 [0.68]	1.89 [0.99]	1.72 [0.94]
GI subjects	1.63 [0.54]	1.71 [0.82]	1.70 [0.72]
Lipid clinic subjects	1.71 [0.71]	1.82 [0.74]	1.80 [0.83]
All subjects	1.67 [0.35]	1.76 [0.78]	1.75 [0.77]

Table 5: Mean changes from baseline at 8 and 16 weeks [mmol/L] and 95% confidence intervals [all patients, n = 63]

	8 wks	p	16 wks	p
TOTAL CHOLESTEROL	0.080 [-0.061,0.22]	p = 0.26	0.068 [-0.056,0.19]	p = 0.28
LDL CHOLESTEROL	0.002 [-0.13,0.14]	p = 0.98	0.018 [-0.098,0.13]	p = 0.75
HDL CHOLESTEROL	0.028 [-0.023,0.08]	p = 0.28	0.012 [-0.035,0.059]	p = 0.61
TRIGLYCERIDES	0.094 [-0.059,0.25]	p = 0.22	0.082 [-0.050,0.21]	p = 0.22

Table 6: Weight and Dietary Intake [SD] at Baseline and at 16 Weeks

	Baseline	16 wks	p
WEIGHT [Kg]	74.6 [16.5]	75.4 [16.9]	p = 0.0039
ENERGY [kcal]	1649 [409]	1689 [419]	p = 0.29
FAT [g]	49.1 [20.3]	49.8 [20.2]	p = 0.66
FIBER [g]	19.7 [7.96]	19.7 [7.28]	p = 0.99

The mean caloric intake or the intakes of fat or fibre did not change significantly during the study [table 6]. There was a small but significant increase in the weights of subjects, which was felt to reflect the high proportion of subjects beginning the study in the fall and the associated reduction of physical activity during the subsequent winter months. There was no apparent relationship between change in cholesterol and change in weight [correlation = .1, p = .15].

Discussion

We failed to prove our hypothesis that administration of psyllium in the morning would have a greater cholesterol-lowering effect than it would in the evening. Not only was there no observable difference in lipid levels between the

crossover periods but the daily ingestion of a greater daily dose than the 10.2 g of psyllium for which the FDA allows health claims to be made [6] had no effect on lipid levels in our study group. No change in any lipid parameter, including total and LDL cholesterol was observed. No difference was found when subgroup analysis was undertaken for the sex of the patients, the time of day they took their psyllium, or whether they were recruited from the gastroenterology clinic or the lipid clinic.

We used a crossover design since this was the most appropriate one for the primary question being addressed; accordingly, our study did not include a control group. However, the nature of the study should not have provided any motivation for study subjects to adopt any new

lifestyle or dietary changes beyond those implemented well before the introduction of psyllium. Observational data has been shown to provide valid information, which is consistent with that observed in randomized, controlled trials [17,18]. The nature of the intervention was in keeping with those undertaken in day-to-day clinical practice and the protocol used should, therefore, have high "clinical relevance".

Failure of lipid-lowering by psyllium has also been demonstrated in twenty hypercholesterolemic children [19], in twenty-four hyperlipidemic adults [20] and in a large observational study of elderly patients taking psyllium [21]. A report of lipid-lowering therapies in hypercholesterolemic veterans showed only a 2% reduction in LDL cholesterol and a small increase in LDL/HDL ratio in patients taking psyllium, but does not provide a measure of statistical significance [22]. One study revealed no difference in total cholesterol-lowering compared to placebo, but a reduction of LDL cholesterol resulted from psyllium treatment [23]. Another demonstrated no difference in total cholesterol-lowering compared to placebo, a reduction of LDL in 11 "responders" and no change in 9 "nonresponders" [24]. A reduction of HDL cholesterol has been noted in some studies [25-27] and was associated with changes in LDL/HDL ratios similar to placebo treatment [25,26].

Published studies include few normocholesterolemic subjects. Cholesterol reduction was observed in 7 normal men [28] and in 5 of 9 subjects [29], in both studies after 3 weeks of treatment. A reduction of cholesterol levels was also observed in 12 elderly patients given psyllium for 4 months [30], while 5 normocholesterolemic subjects in another study showed no reduction after 2 to 7 months of treatment [31].

A meta-analysis of 17 studies of patients with hypercholesterolemia has suggested a small but significant cholesterol-lowering effect of psyllium [2]. All of these investigations were associated in one way or another with the product manufacturer. Additional studies have also indicated some cholesterol-lowering by psyllium in hypercholesterolemic individuals [32-37] or in diabetics [38-40]; however, much of this work is uncontrolled and some protocols have specifically excluded premenopausal women [33,38]. The association of cholesterol-lowering effects with psyllium may be weakened in some studies by the use of a supplement containing additional forms of soluble fibre [42] or by apparent differences in intake of calories [43-46], soluble fibre [25] or cholesterol [47] in control and treatment groups or periods. Several reports include only small numbers of patients and/or are of short duration. There is a strong predominance of male

subjects in these publications and some protocols incorporate additional treatment interventions [20,48].

Several factors may contribute to the difference between our observations and those of others. A meta-analysis has demonstrated that the initial level of cholesterol was highly predictive of the subsequent reduction of cholesterol by oat bran [49]. A greater effect of psyllium in men compared to women has been suggested [23,46] and a diet high in soluble fibre produced less cholesterol-lowering in post menopausal women than in men [10]. Soluble fibre has a lesser effect on lipid metabolism in female than in male guinea pigs [50] and there is a sex-based difference in mechanism of action in this animal [51]. Oat bran fails to lower cholesterol in young women, in contrast to men and older women [52]. The dominance of women in our study, the "normal", or only slightly abnormal cholesterol states of our subjects and the relatively young ages of some of them may, accordingly, account for some of the variance of our observations with some of those previously reported.

The small increase in the weight of subjects is believed to be have resulted from reduced physical activity. In a meta-analysis of the effect of weight reduction on lipids, predominantly through dietary change, a reduction in total cholesterol of 0.05 mmol/L and of 0.02 mmol/L in LDL cholesterol per kilogram of weight lost was identified [53]. Dietary intakes were stable throughout our study and the average weight gain of less than one kilogram is very unlikely to have raised cholesterol levels to a degree sufficient to offset a significant cholesterol lowering effect of psyllium.

A small cholesterol-lowering effect of psyllium appears to occur in hypercholesterolemic individuals, at least in men and possibly postmenopausal women. The notion of a benefit accruing to the general population requires additional study. The promotion of foods containing psyllium as reducing the risk of heart disease for the population at large [6] may be premature. Additional study is required and this should be undertaken in a manner that is free from concern regarding the possibility of publication bias which Brown L, Rosner B, Willett WW and Sacks FM have raised [2].

Conclusion

The timing of psyllium administration had no effect on cholesterol-lowering and, in fact, no cholesterol-lowering was observed. Conclusions regarding the effectiveness of psyllium for the prevention of heart disease in the population at large may be premature.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

GVR carried out the study design, data review and writing. EAS carried out the study design and data review. RB carried out the study design, data review and statistical analysis. ALE carried out the study design and data review.

Acknowledgements

We are indebted to Gwendolyn Shah and Sandra Sharkey for their assistance with the interviews, instruction of subjects and with the gathering of the data and to Victoria Stagg for her help with the statistical analysis.

References

- Anderson JW, Deakins DA, Bridges SR: **Soluble Fibre – Hypocholesterolemic Effects and Proposed Mechanisms.** In: *Dietary Fibre – Chemistry, Physiology and Health Effects* Edited by: Krichevsky D, Bonfield C, Anderson JW. New York, NY: Plenum Press; 1990:339-363.
- Brown L, Rosner B, Willett WW, Sacks FM: **Cholesterol-lowering effects of dietary fibre: a meta-analysis.** *Am J Clin Nutr* 1999, **69**:30-42.
- Hopewell RD, Yeater R, Ulluich I: **Soluble Fibre: Effect on Carbohydrate and Lipid Metabolism.** *Progress In Food and Nutritional Science* 1993, **17**:159-182.
- Lipsky H, Gloger M, Frishman WH: **Dietary Fibre for Reducing Blood Cholesterol.** *Clin Pharmacol* 1990, **30**:699-703.
- Truswell AS: **Dietary fibre and blood lipids.** *Current Opinion in Lipidology* 1995, **6**:14-19.
- Federal Register* 1998, **63**:8103-8121.
- Chaplin MF, Chaudhury S, Dettmar PW, Sykes J, Shaw AD, Davies GJ: **Effect of ispaghula husk on the faecal output of bile acids in healthy volunteers.** *J Biochem Molec Biol* 2000, **72**:283-292.
- Fernandez ML: **Distinct mechanisms of plasma LDL lowering by dietary fibre in the guinea pig: specific effects of pectin, guar gum, and psyllium.** *J Lipid Res* 1995, **36**:2394-2404.
- Forman DT, Garvin JE, Forestiner JE, Taylor CB: **Increased Excretion of Faecal Bile Acids by an Oral Hydrophilic Colloid.** *Proc Soc Exper Biol Med* 1968, **127**:1060-1063.
- Jenkins DJ, Wolever TM, Rao AV, Hegele RA, Mitchell SJ, Ransom TP, Boctor DL, Spadafora PJ, Jenkins AL, Mehling C: **Effect on blood lipids of very high intakes of fibre in diets low in saturated fat and cholesterol.** *NEJM* 1993, **329**:21-6.
- Kesaniemi YA, Tarpila S, Miettinen TA: **Low vs. high dietary fibre and serum, biliary, and faecal lipids in middle-aged men.** *Am J Clin Nutr* 1990, **51**:1007-12.
- Romero AL, West KL, Zern T, Fernandez ML: **The seeds from Plantago ovata lower plasma lipids by altering hepatic and bile acid metabolism in guinea pigs.** *J Nutr* 2002, **132**:1194-8.
- Stanley MM, Paul D, Gacke D, Murphy J: **Effects of Cholestyramine, Metamucil, and Cellulose on Faecal Bile Salt Excretion in Man.** *Gastroenterology* 1973, **65**:889-894.
- Turley SD, Daggy BP, Dietschy JM: **Cholesterol-Lowering Action of Psyllium Mucilloid in the Hamster: Sites and Possible Mechanisms of Action.** *Metabolism* 1991, **40**:1063-1073.
- Vahouny GV, Tombes R, Cassidy MM, Krichevsky D: **Gallo LL, Dietary Fibres: V. Binding of Bile Salts, Phospholipids and Cholesterol from Mixed Micelles by Bile Acid Sequestrants and Dietary Fibres.** *Lipids* 1980, **15**:1012-1018.
- Ho K, Drummond JL: **Circadian rhythm of biliary excretion and its control mechanisms in rats with chronic biliary drainage.** *Am J Physiol* 1975, **229**:1427-1437.
- Benson K, Hartz AJ: **A Comparison of Observational Studies and Randomized, Controlled Trials.** *N Eng J Med* 2000, **342**:1878-86.
- Concato J, Shah N, Horwitz RL: **Randomized, Controlled Trials, Observational Studies, and the Hierarchy of Research Designs.** *N Eng J Med* 2000, **342**:1887-92.
- Dennison BA, Levine DM: **Randomized, double-blind, placebo-controlled, two-period crossover clinical trial of psyllium fibre in children with hypercholesterolemia.** *J Pediatr* 1993, **123**:24-29.
- Spence JD, Huff MW, Heidenheim P, Viswanatha A, Munoz C, Lindsay R, Wolfe B, Mills D: **Combination Therapy with Colestipol and Psyllium Mucilloid in patients with Hyperlipidemia.** *Ann Intern Med* 1995, **123**:493-499.
- Stewart RB, Hale WE, Moore MT, May FE: **Marks RG, Effect of Psyllium Hydrophilic Mucilloid on Serum Cholesterol in the Elderly.** *Dig Dis Sci* 1991, **36**:329-334.
- Schectman G, Hiatt J, Hartz A: **Evaluation of the Effectiveness of Lipid Lowering Therapy (Bile Acid Sequestrants, Niacin, Psyllium and Lovastatin) for Treating Hypercholesterolemia in Veterans.** *Am J of Cardiol* 1993, **71**:759-765.
- Anderson JW, Floore TL, Geil PB: **Hypocholesterolemic Effects of Different Bulk-Forming Hydrophilic Fibres as Adjuncts to Dietary Therapy in Mild to Moderate Hypercholesterolemia.** *Intern Med* 1991, **151**:1597-1602.
- Everson GT, Daggy BP, McKinley C, Story JA: **Effects of psyllium hydrophilic mucilloid on LDL-cholesterol and bile acid synthesis in hypercholesterolemic men.** *J Lipid Res* 1992, **33**:1183-1192.
- Levin EG, Miller VT, Muesing RA, Stoy DB, Balm TK, LaRosa JC: **Comparison of Psyllium Hydrophilic Mucilloid and Cellulose as Adjuncts to a Prudent Diet in the Treatment of Mild to Moderate Hypercholesterolemia.** *Arch Intern Med* 1990, **150**:1822-1827.
- Wolever TMS, Jenkins DJA, Mueller S, Boctor DL, Ransom TP, Patten R, Chao ES, McMillan K, Fulgoni V: **Method of administration influences the serum cholesterol-lowering effect of psyllium.** *Am J Clin Nutr* 1994, **59**:1055-1059.
- Wolever TMS, Jenkins DJA, Mueller S, Patten R, Relle LK, Boctor D, Relle LK, Ransom TP, Chao ES, McMillan K, Fulgoni V: **Psyllium Reduces Blood Lipids in Men and Women With Hyperlipidemia.** *Am J Med Sci* 1994, **307**:269-273.
- Abraham ZD, Mehta T: **Three-week psyllium-husk supplementation: effect on plasma cholesterol concentrations, faecal steroid excretion, and carbohydrate absorption in men.** *Am J Clin Nutr* 1988, **47**:67-74.
- Lieberthal MM, Martens RA: **Lowered Serum Cholesterol Following the Ingestion of a Hydrophilic Colloid.** *AM J Dig Dis* 1975, **20**:469-474.
- Burton R, Manninen V: **Influence of a Psyllium-based Fibre Preparation on Faecal and Serum Parameters.** *Acta Med Scand (Suppl)* 1982, **668**:91-94.
- Danielsson A, Ek B, Nyhlin H, Steen L: **Effect of Long Term Treatment with Hydrophilic Colloid on Serum Lipids.** *Acta Hepato-Gastroenterol* 1979, **26**:148-153.
- Davidson MH, Maki KC, Kong JC, Dugan LD, Torri SA, Hall HA, Drennan KB, Anderson SM, Fulgoni VL, Saldanha LG, Olson BH: **Long-term effects of consuming foods containing psyllium seed husk on serum lipids in subjects with hypercholesterolemia.** *Am J Clin Nutr* 1998, **67**:367-76.
- Jenkins DJA, Wolever TMS, Vidgen E, Kendall CW, Ransom TP, Mehling CC, Mueller S, Cunnane SC, O'Connell NC, Setchell KD, Lau H, Teitel JM, Garvey MB, Fulgoni V 3rd, Connelly PW, Patten R, Corey PN: **Effect of psyllium in hypercholesterolemia at two monounsaturated fatty acid intakes.** *Am J Clin Nutr* 1997, **65**:1524-33.
- MacMahon M, Carless J: **Ispaghula husk in the treatment of hypercholesterolaemia: a double-blind controlled study.** *J Cardio Vasc Risk* 1998, **5**:167-172.
- Romero AL, Romero JE, Galaviz S, Fernandez ML: **Cookies Enriched with Psyllium or Oat Bran Lower Plasma LDL Cholesterol in Normal and Hypercholesterolemic Men from Northern Mexico.** *J Am Coll Nutr* 1998, **17**:601-608.
- Segawa K, Kataoka T, Fukuo Y: **Cholesterol-Lowering Effects of Psyllium Seed Associated with Urea Metabolism.** *Biol Pharm Bull* 1998, **21**:184-187.
- Weingand KW, Le N, Kuzmak BR, Brown WV, Daggy BP, Miettinen TA, Howard BV, Howard WJ: **Effects of Psyllium on Cholesterol and Low-density Lipoprotein Metabolism in Subjects with Hypercholesterolemia.** *Endocrinol Metab* 1997, **4**:141-150.
- Fagerberg S: **The effects of a bulk laxative (Metamucil) on fasting blood glucose, serum, lipids and other variables in constipated patients with non-insulin dependent adult diabetes.** *Curr Ther Res* 1982, **31**:166-172.
- Fрати-Munari AC, Fernandez-Harp JA, Becerril M, Chavez-Negrete A, Banales-Ham M.: **Decrease in serum lipids, glycemia and body-weight by plantago psyllium in obese and diabetic patients.** *Arch Invest Med (Mex)* 1983, **14**:259-268.

40. Gupta RR, Agrawal CG, Singh GP, Ghatak A: **Lipid-lowering efficacy of psyllium hydrophilic mucilloid in non insulin dependent diabetes mellitus with hyperlipidaemia.** *Indian J Med Res* 1994, **100**:237-241.
41. Jenkins DJ, Kendall CW, Vuksan V, Vidgen E, Parker T, Faulkner D, Mehling CC, Garsetti M, Testolin G, Cunnane SC, Ryan MA, Corey PN: **Soluble fibre intake at a dose approved by the US Food and Drug Administration for a claim of health benefits: serum lipid risk factors for cardiovascular disease assessed in a randomized controlled crossover trial.** *Am J Clin Nutr* 2002, **75**:834-9.
42. Roberts DCK, Truswell AS, Bencke A, Dewar HM, Farmakalidis E: **The cholesterol-lowering effect of a breakfast cereal containing psyllium fibre.** *Med J Aust* 1994, **161**:660-664.
43. Anderson JW, Riddell-Mason S, Gustafson NJ, Smith SF, Mackey M: **Cholesterol-lowering effects of psyllium-enriched cereal as an adjunct to a prudent diet in the treatment of mild to moderate hypercholesterolemia.** *Am J Clin Nutr* 1992, **56**:93-98.
44. Bell LP, Hectorn KJ, Reynolds H, Balm TK, Hunninghake DB: **Cholesterol-Lowering Effects of Psyllium Hydrophilic Mucilloid: Adjunct Therapy to a Prudent Diet for Patients With Mild to Moderate Hypercholesterolemia.** *JAMA* 1989, **261**:3419-3423.
45. Bell LP, Hectorn KJ, Reynolds H, Hunninghake DB: **Cholesterol-lowering effects of soluble-fibre cereals as part of a prudent diet for patients with mild to moderate hypercholesterolemia.** *Am J Clin Nutr* 1990, **52**:1020-1026.
46. Vega-Lopez S, Conde-Knape K, Vidal-Quintanar RL, Shachter NS, Fernandez ML: **Sex and hormonal status influence the effects of psyllium on lipoprotein remodelling and composition.** *Metabolism* 2002, **51**:500-7.
47. Sprecher DL, Harris BV, Goldberg AC, Anderson EC, Bayuk LM, Russell BS, Crone DS, Quinn C, Bateman J, Kuzmak BR: **Efficacy of Psyllium in Reducing Serum Cholesterol Levels in Hypercholesterolemic Patients on High-or Low-Fat Diets.** *Ann Intern Med* 1993, **119**:545-554.
48. Maciejko JJ, Brazg R, Shah A, Patil S, Rubenfire M: **Psyllium for the Reduction of Cholestyramine-Associated Gastrointestinal Symptoms in the Treatment of Primary Hypercholesterolemia.** *Arch Fam Med* 1994, **3**:955-960.
49. Ripsin CM, Keenan JM, Jacobs DR Jr, Elmer PJ, Welch RR, Van Horn L, Liu K, Turnbull WH, Thye FW, Kestin M: **Oat Products and Lipid Lowering: A Meta-analysis.** *JAMA* 1992, **267**:3317-3325.
50. Fernandez ML, Vergara-Jimenez M, Romero AL, Erickson SK, McNamara DJ: **Gender differences in response to dietary soluble fibre in guinea pigs: effects of pectin, guar gum, and psyllium.** *J Lipid Res* 1995, **36**:2191-2202.
51. Shen H, He L, Price RL, Fernandez ML: **Dietary soluble fibre lowers plasma LDL cholesterol concentrations by altering lipoprotein metabolism in female guinea pigs.** *J Nutr* 1998, **128**:1434-41.
52. Keenan JM, Wenz JB, Myers S, Ripsin C, Huang ZQ: **Randomized, Controlled, Crossover Trial of Oat Bran in Hypercholesterolemic Subjects.** *J Fam Pract* 1991, **33**:600-608.
53. Dattilo AM, Kris-Etherton PM: **Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis.** *Am J Clin Nutr* 1992, **56**:320-8.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

