

# **Soy foods have low glycemic and insulin response indices in normal weight subjects**

Robert M. Blair, Ph.D.\*, E.C. Henley, Ph.D., Aaron Tabor, M.D.

Physicians Pharmaceuticals, Inc., 1031 E. Mountain St., Building 302, Kernersville,  
NC 27284

\*Corresponding author

Email addresses:

RMB: [drblair@revivalsoy.com](mailto:drblair@revivalsoy.com)

ECH: [henleyec@aol.com](mailto:henleyec@aol.com)

AT: [drtabor@revivalsoy.com](mailto:drtabor@revivalsoy.com)

Corresponding Author:

Robert M. Blair, Ph.D.

Research Manager

Physicians Pharmaceuticals, Inc.

1031 E. Mountain St., Building 302

Kernersville, NC 27284

Telephone: 336-722-2337, x1734

Fax: 336-722-7712

# **Abstract**

## **Background**

Foods with a low glycemic index (GI) may provide a variety of health benefits. The objective of the present study was to measure the GI and insulin index (II) of select soy foods.

## **Methods**

The study was conducted in two parts with low-carbohydrate products being tested separately. In Experiment 1, subjects averaged 23.2 years of age with BMI = 22.0 kg/m<sup>2</sup>, while subjects in Experiment 2 averaged 23.9 years of age with BMI = 21.6 kg/m<sup>2</sup>. The reference (glucose) and test foods were served in portions containing 10 g of carbohydrates in Experiment 1 and 25 g of carbohydrates in Experiment 2.

Subjects consumed the reference food twice and each test food once. For each test, subjects were instructed to consume a fixed portion of the reference food or test food together with 250 g of water within 12 min. Blood samples were collected before each test and at 15, 30, 45, 60, 90, and 120 min after consumption of reference or test foods to quantify glucose and insulin. Two-hour blood glucose and plasma insulin curves were constructed and areas under the curves were calculated. GI and II values for each subject and test food were calculated. Analysis of variance and the Fisher PLSD test for multiple comparisons were used to test for significant differences between the test foods' mean GI and II values.

## **Results**

In Experiment 1, both low-carbohydrate soy foods were shown to have significantly ( $P < 0.05$ ) lower GI and II values than the reference food. In Experiment 2, three of

the four test foods had significantly ( $P < 0.05$ ) lower GI and II values than the reference food.

### **Conclusions**

All but one of the soy foods tested had a low GI, suggesting that soy foods may be an appropriate part of diets intended to improve control of blood glucose and insulin levels.

## Background

The glycemic index (GI) was first developed by Jenkins and colleagues (1) as a new method of classifying foods based on the blood glucose response after food consumption. The GI value of a food is a percentage of the 2-hour area under the blood glucose response curve of a reference food, typically glucose (2). Since the GI is determined for a particular quantity of carbohydrates in the food being tested and since the actual amount of carbohydrates consumed in a meal or snack varies greatly, the GI concept was expanded to include the concept of glycemic load (GL). The GL is determined by multiplying the GI of a food by the grams of carbohydrates in a serving. The GL value incorporates the amount of digestible carbohydrates in a serving in order to better gauge the impact of a meal or snack on postprandial glucose response (3, 4).

It has been reported that a high GI diet may have adverse health consequences by increasing the risk for chronic disease (5, 6). Recent evidence suggests that high GI/GL diets may increase the risk for cardiovascular disease (CVD) (7-9) and type 2 diabetes (3, 4, 10, 11). A high GI diet may increase the risk for chronic disease through the stimulation of hyperglycemia and hyperinsulinemia (6).

In contrast, a low GI diet has been reported to have health benefits (5, 6, 12, 13). A low GI diet has been shown to improve glycemic control (14-17), aid in weight loss (18, 19), and reduce some CVD risk factors (9, 14, 20-23).

To date, only about 30 – 40 soy foods have been assessed for their GI/GL values (1, 24-27). The objective of the current study was to determine the GI and insulin index (II) values of select soy food products (bars, drinks, pasta, and chips) currently available on the U.S. and international markets.

## **Methods**

The current study was conducted using internationally recognized GI methodology (28-30). The experimental procedures used in this study were in accordance with international standards for conducting ethical research with humans and were approved by the Human Research Ethics Committee of Sydney University of Australia where the study was conducted by contract.

A search of the literature using the National Library of Medicine's PubMed search engine showed a paucity of literature on the determination of the glycemic index of soy foods. A search of "soy AND glycemic index" yielded 5 hits; a search of "soy AND glycemic load" yielded 2 hits; a search of "soy AND glycemic" yielded 14 hits and a search of "isoflavone AND glycemic" yielded 5 hits. However, none of the papers found measured the glycemic index or glycemic load of soy foods. A search of the international table of glycemic index and glycemic load values (26) showed that the glycemic index of several soy foods have been measured, but only a few of these have been reported as independent publications.

## **Study Subjects**

For both experiments, 10 healthy, non-smoking subjects, were recruited from the staff and student population of the University of Sydney. Exclusion criteria included being overweight, dieting, impaired glucose tolerance, illness or food allergy, or regular use of prescription medication (other than contraceptive medication). All study participants gave written informed consent before participating in this study.

In Experiment 1 the 10 subjects (two females, eight males) had a mean age of 23.2 years (19.9–25.7 years) and a mean body mass index (BMI) score of 22.0 kg/m<sup>2</sup> (absolute range = 19.4–25.4 kg/m<sup>2</sup>). In Experiment 2 the 10 subjects (four females, six males) had a mean age of 23.9 years (20.3–26.9 years) and a mean BMI score of 21.6 kg/m<sup>2</sup> (absolute range = 19.5–25.8 kg/m<sup>2</sup>). Four subjects participated in both experiments.

### **Composition of Test Foods**

In Experiment 1, the reference and two low-carbohydrate test foods (products with  $\leq 6$  g net carbohydrates (available carbohydrates); net carbohydrates = total carbohydrates – sugar alcohols – fiber) were served to subjects in portions containing 10 g of digestible carbohydrate. In Experiment 2, the reference and four test foods were served to subjects in portions containing 25 g of available carbohydrate. Glucose (Glucodin® powder, Boots Health Care Company, North Ryde, NSW Australia) dissolved in water was the reference food. Mass and nutrient contents of the reference and test foods (Revival Soy® from Physicians Pharmaceuticals, Inc., Kernersville, NC, USA) are listed in **Tables 1 and 2**.

Each portion of the reference food was prepared the day before required by dissolving the glucose in 250 g of water and storing overnight at 4 °C. The individual portions of the test foods were prepared the day before required, except for the soy spaghetti. The individual portions of the uncooked soy spaghetti were weighed the day before required. On the testing day, each portion of dry spaghetti was cooked for 4 minutes in boiling water and drained. The reference and test foods were served with 250 g of plain water. The subjects consumed all the food and fluid served to them at a comfortable pace within 12 minutes.

### **Experimental Procedures**

The experimental methods used in the current study have been previously described (31) and are briefly outlined here. In both experiments, study subjects consumed the reference food on two separate occasions and each of the test foods on one occasion only after a 10-hour overnight fast. The reference food was consumed on the first and last test sessions, and test foods were consumed in random order in between. Each test session was completed on a separate morning with at least a day between subsequent sessions.

On each test day a baseline, finger-prick blood sample was obtained for blood glucose and plasma insulin determinations using an automatic, non-reusable lancet device (Safe-T-Pro®, Boehringer Mannheim GmbH, Mannheim, Germany). Following consumption of the reference or test food, additional blood samples were collected at 15, 30, 45, 60, 90 and 120 minutes. Blood glucose concentrations were measured

immediately after the blood samples were collected. Blood samples collected for plasma insulin determination were centrifuged for 30 seconds immediately after collection and the plasma layer from each sample was transferred into a labeled, uncoated microcentrifuge tube and stored at -20 °C until analyzed.

### **Blood Glucose and Glycemic Index Determinations**

The glucose concentration in each of the whole capillary blood samples was analyzed in duplicate using a glucose dehydrogenase/mutarotase enzymatic reaction using a HemoCue® beta-glucose photometric analyzer (HemoCue AB, Ängelholm, Sweden). Duplicate readings were accepted if the two separate measurements for each timepoint were within 0.3 mmol/L of each other. If the readings were not within 0.3 mmol/L of each other, than an additional 2 blood glucose sample readings were taken from the subject within approximately 40 – 60 seconds after the initial readings. The two or three similar (i.e. within 0.3 mmol/L) readings were then averaged together to obtain the blood glucose response for that time point. A two-hour blood glucose response curve was constructed and the incremental area under the glucose response curve (IAUC) was calculated. The GI value for each test food was calculated for each subject by dividing the two-hour blood glucose IAUC value for the test foods by their average two-hour blood glucose IAUC value for the reference food and multiplying by 100 to obtain a percentage score. The final reported GI value for each test food is the mean GI value for that food in the group of 10 subjects.

### **Plasma Insulin and Insulin Index Determinations**

The concentration of insulin in each plasma sample was analyzed using a solid-phase antibody-coated tube radioimmunoassay kit (Diagnostic Products Corporation, Los Angeles, CA, USA). A two-hour plasma insulin response curve was constructed from the data and the IAUC of the insulin response curve was calculated. The II value for each test food was calculated for each subject by dividing their plasma insulin IAUC value for the test foods by their mean plasma insulin IAUC value for the reference food and multiplying by 100 to obtain a percentage score.

### **Statistical Analyses**

Sample size calculations (90% power, level of significance = 0.05) using data from published GI studies indicated that a minimum of eight study subjects would be needed to detect significant differences among the GI values of the reference and test foods. Analysis of variance (ANOVA) and the Fisher PLSD test for multiple comparisons were used to determine significant differences between the test foods' mean GI and II values. Statistical analyses were conducted using Statview Student<sup>TM</sup> Software (version 4, Abacus Concepts Inc., Berkley, CA, USA). Significance was assumed at  $P < 0.05$ .

## **Results**

### **Experiment 1: Low-Carbohydrate Soy Foods**

The mean blood glucose response curves for the reference and two test foods are shown in **Figure 1**. The reference food produced a much larger rise in blood glucose during the first 30 minutes and a greater overall glycemic response than the two test foods. The two test foods produced slightly different glucose response curves with the Chocolate Raspberry Zing™ bar producing a higher glycemic response than the Chocolate Daydream™ sucralose shake. However, both foods produced very low glycemic response curves.

The mean plasma insulin response curves for the reference and two test foods are shown in **Figure 2**. The plasma insulin responses observed for the reference food and the test foods showed a similar profile to their concurrent blood glucose responses. The reference food produced the highest peak plasma insulin concentration and the largest overall plasma insulin response, followed by the Chocolate Raspberry Zing™ bar and the Chocolate Daydream™ sucralose shake, respectively.

## **Experiment 2: Other Soy Foods**

The mean blood glucose response curves for the reference and the four test foods are shown in **Figure 3**. Similar to the results observed in Experiment 1, the reference food produced a large rise in blood glucose during the first 30 minutes and the greatest overall glycemic response. The four test foods varied in their peak blood glucose concentrations and their overall glycemic responses. Among the test foods, the soy pasta chips produced the largest glycemic response followed by the Peanut Butter Chocolate Pal™ bar, the soy spaghetti, and the Chocolate Daydream™ fructose shake.

The mean plasma insulin response curves for the reference and the four test products are shown in **Figure 4**. The foods' average plasma insulin responses were similar to their respective mean plasma glucose responses. The reference food produced the largest plasma insulin response, followed by the four test foods in the same order as their glycemic responses.

### **Glycemic and Insulin Indices**

The mean GI value of the glucose reference was significantly greater ( $P < 0.001$ ) than the mean GI values of each of the test foods with the exception of the baked soy pasta chips (**Figure 5**). The mean GI value of the soy pasta chips was not different from the glucose reference, but was significantly greater ( $P < 0.001$ ) than the mean GI values for the other five test foods. Despite a high GI value, the soy pasta chips had only a medium GL value due to the small serving size and relatively low carbohydrate level (**Table 3**).

The mean II value of the glucose reference was significantly greater ( $P < 0.001$ ) than the mean II values of each of the six test foods (**Figure 5**). The mean II of the soy pasta chips was significantly higher ( $P < 0.001$ ) than the mean II values for the soy spaghetti, Chocolate Daydream™ fructose shake, Chocolate Raspberry Zing™ bar, and the Chocolate Daydream™ sucralose shake. The mean II values for the Peanut Butter Chocolate Pal™ bar and the Chocolate Raspberry Zing™ bar were both significantly greater ( $P < 0.001$ ) than the mean II values for the Chocolate Daydream™ fructose and Chocolate Daydream™ sucralose shakes.

## Discussion

The results of this study demonstrate that 5 of the 6 soy food products tested have a low GI value ( $GI \leq 55$ ). Of the 6 products tested, only the baked soy pasta chips had a high GI value ( $GI > 70$ ). However, when the amount of available carbohydrates in one serving of the soy pasta chips was used to calculate a GL value, the soy pasta chips had a medium GL value. The other products tested also were either low or medium GL foods.

An increasing body of evidence suggests that the GI and/or GL values of foods impact human health (see 5, 6, 12, 32, and 33 for reviews). Low-GI diets have been shown to improve glycemic control in diabetic (14, 15) and non-diabetic (16, 17) subjects. Diet impacts the incidence of type 2 diabetes and the regulation of dietary carbohydrate has taken on a prominent role in dietary control of this chronic disease. Two recent meta-analyses (13, 34) reported that consumption of low-GI foods rather than high-GI foods appears to modestly improve glycemic control by reducing plasma cholesterol, fructosamine, and hemoglobin A<sub>1c</sub> (Hb<sub>A1c</sub>) levels.

A number of studies suggest that high GI/GL diets may increase CVD risk (7-9), while several others indicate that low GI diets may reduce some CVD risk factors (9, 14, 15, 20-23). Meta-analysis results indicated that low GI diets significantly reduced total cholesterol (average reduction = 0.17 mmol/L;  $P = 0.03$ ) and HbA<sub>1c</sub> (average reduction after 12 weeks = 0.45%;  $P = 0.02$ ) compared to high GI diets (35). These

data suggest that dietary GI may improve some, but not all markers of cardiovascular disease risk.

Few studies have examined the effect of low GI diets on weight loss; however, there is some evidence that low GI diets may be beneficial (33). In obese women, an energy-restricted, low GI diet significantly increased weight loss compared to an energy-restricted, high GI diet (18). Spieth and co-workers (19) demonstrated that an *ad libitum* low GI diet significantly ( $P < 0.05$ ) reduced BMI to a greater extent than did an energy-restricted, low fat diet. A recent study demonstrated that dietary GI was inversely associated with thigh intramuscular fat while GL was inversely associated with visceral abdominal fat in men (36). Despite the potential benefits of low-GI diets on weight loss, a number of studies report no effect on weight loss (37, 38).

Soy protein is a high quality protein that has been extensively studied. The quality of soy protein has been assessed through several metabolic studies of nitrogen balance (39-41), which have demonstrated that soy protein supports nitrogen balance on par with beef and milk proteins. One recent study reported that amino acids from soy protein appear in the serum sooner, but that this may lead to a more rapid breakdown of the amino acids in the liver (42).

Dietary soy consumption has been shown to have beneficial effects on several aspects of human health, including the diseases potentially influenced by dietary GI levels (43-45). Soy consumption has been reported to modestly improve plasma lipid profiles (46, 47), improve bone health (48), help reduce menopausal symptoms (49),

and slightly reduce the risk of breast (50) and prostate cancers (51). The health benefits of dietary soy have been attributed to its isoflavones as well as to the biological actions of its constituent proteins. However, an additional means of providing health benefits may be through the low GI of soy and soy foods.

The international table of GI and GL values (26) reports the GI/GL values of a number of soy foods. These values range from a low GI of 14 for soybeans canned in brine to a high GI of 115 for a tofu-based frozen dessert (24). The Revival Soy® products tested in this study fell within this range and with the exception of the baked soy pasta chips were all within the low GI category. Only a few other studies have reported on the GI of soy-based foods. Packer et al. (25) indicated that gluten-free, soy-based, bread had a high GI value. In contrast, the addition of soy foods has been shown to lower the GI value of mixed meals (27). Similarly, the replacement of unrefined wheat flour with soy flour lowers the GI value of parantha, an Indian snack food (52). Similar to the Revival Soy® bars tested in the current study, other snack bars containing soy have been shown to have low to medium GI values (53). Previous data report that the GI values of spaghetti ranges from 27 – 68. The Revival Soy® thin spaghetti had a GI value = 47, similar to other spaghetti products. Overall, these studies indicate that soy-based foods generally have a low to medium GI value and would be suitable for individuals concerned with regulating blood glucose and insulin levels.

The ingredients and form of a food product affect its GI value. For example, while soybeans have a low GI value, the use of high GI ingredients in soy foods can increase the GI value of the final product. This was likely the case with the baked soy

pasta chips. The baked soy pasta chips contain potato starch and potatoes have a high GI value (33). Additionally, the baked soy pasta chips have a puffed physical form, which may lead to high GI values. Similar to planning diets with types of fat and protein in mind, types of carbohydrates should also be considered since carbohydrate types influence the GI. The substitution of high GI ingredients with low GI ingredients in food products like the baked soy pasta chips may help keep the final GI value down.

In addition to the effects of form and content of foods on the GI value, consumption of other foods with low GI foods can affect the overall GI value of meal. Sugiyama et al. (34) demonstrated that adding soybean products (miso, natto, and ground soybean) lowered the GI of white rice by 20 – 40%. However, further studies are required before any conclusions can be drawn.

## **Conclusions**

With the apparent resurgence of interest in low-GI diets for weight loss and health benefits, it is important that information on the GI value of foods is available. Therefore, we conducted the current study to determine the GI value of a small variety of commercially available soy foods. The results of the current study demonstrate that soy food products generally have low GI values and low to medium GL values. Improvements in ingredient selection and usage may further improve glycemic responses to soy foods. The low GI of soy foods appears to be an additional

benefit of soy for human health and suggests that soy foods are an appropriate part of diet plans intended to improve control over blood glucose and insulin levels.

## **Competing interests**

Dr. Blair is the Research Manager of Physicians Pharmaceuticals, Inc. Dr. Henley is a consultant to Physicians Pharmaceuticals, Inc. Dr. Tabor is the CEO and Medical Research Director of Physicians Pharmaceuticals, Inc.

## **Authors' contributions**

RMB prepared the manuscript. ECH and AT participated in the coordination and design of the study and assisted with manuscript preparation.

## **Acknowledgements**

This study was funded by Physicians Pharmaceuticals, Inc. and conducted via contract research at the University of Sydney, Australia.

## References

- [1] Jenkins DJ, Wolever TM, Taylor RH, Barker H, Fielden H, Baldwin JM, Bowling AC, Newman HC, Jenkins AL: **Glycemic index of foods: a physiological basis for carbohydrate exchange.** *Am J Clin Nutr* 1981, **34**:362-366.
- [2] Wolever TMS, Jenkins DJA, Jenkins AL, Josse RG: **The glycemic index: methodology and clinical implications.** *Am J Clin Nutr* 1991, **54**:846-854.
- [3] Salmeron J, Ascherio A, Rimm EB, Colditz GA, Spiegelman D, Jenkins DJ, Stampfer MJ, Wing AL, Willett WC: **Dietary fiber, glycemic load, and risk of NIDDM in men.** *Diabetes Care* 1997a, **20**:545-50.
- [4] Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willett WC: **Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women.** *JAMA* 1997b, **277**:472-477.
- [5] Augustin LS, Franceschi S, Jenkins DJA, Kendall CWC, La Vecchia C: **Glycemic index in chronic disease: a review.** *Eur J Clin Nutr* 2002, **56**:1049-1071.

- [6] Ludwig DS: **The glycemic index: physiological mechanisms relating to obesity, diabetes, and cardiovascular disease.** *JAMA* 2002, **287**:2414-2423.
- [7] Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, Hennekens CH, Manson JE: **A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women.** *Am J Clin Nutr* 2000, **71**:1455-1461.
- [8] Stampfer MJ, Hu FB, Manson JE, Rimm EB, Willett WC: **Primary prevention of coronary heart disease in women through diet and lifestyle.** *New Engl J Med* 2000, **343**:16-22.
- [9] Amano Y, Kawakubo K, Lee JS, Tang AC, Sugiyama M, Mori K: **Correlation between dietary glycemic index and cardiovascular disease risk factors among Japanese women.** *Eur J Clin Nutr* 2004, **58**:1472-1478.
- [10] Hodge AM, English DR, O'Dea K, Giles GG: **Glycemic index and dietary fiber and the risk of type 2 diabetes.** *Diabetes Care* 2004, **27**:2701-2706.
- [11] Schulze MB, Liu S, Rimm EB, Manson JE, Willett WC, Hu FB: **Glycemic index, glycemic load, and dietary fiber intake and incidence of type 2 diabetes in younger and middle-aged women.** *Am J Clin Nutr* 2004, **80**:348-356.

- [12] Colombani PC: **Glycemic index and load – dynamic dietary guidelines in the context of diseases.** *Physiol Behav* 2004, **83**:603-610.
- [13] Opperman AM, Venter CS, Oosthuizen W, Thompson RL, Vorster HH: **Meta-analysis of the health effects of using the glycaemic index in meal-planning.** *Br J Nutr* 2004, **92**:367-381.
- [14] Jarvi AE, Karlstrom BE, Granfeldt YE, Bjorck IE, Asp NG, Vessby BO: **Improved glycemic control and lipid profile and normalized fibrinolytic activity on a low-glycemic index diet in type 2 diabetic patients.** *Diabetes Care* 1999, **22**:10-18.
- [15] Giacco R, Parillo M, Rivellese AA, Lasorella G, Giacco A, D’Episcopo L, Riccardi G: **Long-term dietary treatment with increased amounts of fiber-rich low-glycemic index natural foods improves blood glucose control and reduces the number of hypoglycemic control in children with type 1 diabetes.** *Diabetes Care* 2000, **23**:1461-1466.
- [16] Jenkins DJ, Wolever TM, Collier GR, Ocana A, Rao AV, Buckley G, Lam Y, Mayer A, Thompson LU: **Metabolic effects of a low-glycemic-index diet.** *Am J Clin Nutr* 1987a, **46**:968-975.
- [17] Frost G, Keogh B, Smith D, Akinsanya K, Leeds S: **The effect of low-glycemic carbohydrate on insulin and glucose response in vivo and in**

- vitro in patients with coronary heart disease.** *Metabolism* 1996, **45**:669-672.
- [18] Slabber M, Barnard HC, Kuyl JM, Dannhauser A, Schall R: **Effects of a low-insulin-response, energy-restricted diet on weight loss and plasma insulin concentrations in hyperinsulinemic obese females.** *Am J Clin Nutr* 1994, **60**:48-53.
- [19] Spieth LE, Harnish JD, Lenders CM, Raezer LB, Pereira MA, Hangen SJ, Ludwig DS: **A low-glycemic index diet in the treatment of pediatric obesity.** *Arch Pediatr Adolesc Med* 2000, **154**:947-951.
- [20] Jenkins DJ, Wolever TM, Kalmusky J, Guidici S, Giordano C, Patten R, Wong GS, Bird JN, Hall M, Buckley G: **Low-glycemic index diet in hyperlipidemia: use of traditional starchy foods.** *Am J Clin Nutr* 1987b, **46**:66-71.
- [21] Luscombe ND, Noakes M, Clifton PM: **Diets high and low in glycemic index versus high monounsaturated fat diets: effects on glucose and lipid metabolism in NIDDM.** *Eur J Clin Nutr* 1999, **53**:473-478.
- [22] Ford ES, Liu S: **Glycemic index and serum high-density lipoprotein cholesterol concentration among US adults.** *Arch Intern Med* 2001, **26**:572-576.

- [23] Liu S, Manson JE, Buring JE, Stampfer MJ, Willett WC, Ridker PM: **Relation between a diet with a high glycemic load and plasma concentrations of high-sensitivity C-reactive protein in middle-aged women.** *Am J Clin Nutr* 2002, **75**:492-498.
- [24] Bukar J, Mezitis NHE, Saitas V, Pi-Sunyer FX: **Frozen desserts and glycaemic response in well-controlled NIDDM patients.** *Diabetes Care* 1990, **13**:382-385.
- [25] Packer SC, Dornhurst A, Frost GS: **The glycaemic index of a range of gluten-free foods.** *Diabet Med* 2000, **17**:657-660.
- [26] Foster-Powell K, Holt SHA, Brand-Miller JC: **International table of glycemic index and glycemic load values: 2002.** *Am J Clin Nutr* 2002, **76**:5-56.
- [27] Sugiyama M, Tang AC, Wakaki Y, Koyama W: **Glycemic index of single and mixed meal foods among common Japanese foods.** *Eur J Clin Nutr* 2003, **57**:743-752.
- [28] Joint FAO/WHO Report. **Carbohydrates in Human Nutrition.** FAO Food and Nutrition, Paper #66. Rome: FAO, 1998.
- [29] Brouns F, Bjorck I, Frayn KN, Gibbs AL, Lang V, Slama G, Wolever TMS: **Glycaemic index methodology.** *Nutr Res Rev* 2005, **18**:145-171.

- [30] Wolever TM, Vorster HH, Bjorck I, Brand-Miller J, Brighenti F, Mann JI, Ramdath DD, Granfeldt Y, Holt S, Perry TL, Venter C, Xiaomei W: **Determination of the glycaemic index of foods: interlaboratory study.** *Eur J Clin Nutr* 2003, **57**:475-482.
- [31] Brand-Miller JC, Thomas M, Swan V, Ahmad ZI, Petocz P, Colagiuri S: **Physiological validation of the concept of glyemic load in lean young adults.** *J Nutr* 2003, **133**:2728-2732.
- [32] Dickinson S, Brand-Miller J: **Glycemic index, postprandial glycemia and cardiovascular disease.** *Curr Opin Lipidol* 2005, **16**:69-75.
- [33] Brand-Miller JC, Holt SHA, Pawlak DB, McMillan J: **Glycemic index and obesity.** *Am J Clin Nutr* 2002, **76(Suppl)**:281S-285S.
- [34] Brand-Miller J, Hayne S, Petocz P, Colagiuri S: **Low-glycemic index diets in the management of diabetes: a meta-analysis of randomized controlled trials.** *Diabetes Care* 2003, **26**:2261-2267.
- [35] Kelly S, Frost G, Whittaker V, Summerbell C: **Low glycaemic index diets for coronary heart disease.** In *The Cochrane Database of Systematic Reviews*. 2004, Issue 4. Art. No.: CD004467.pub2. DOI: 10.1002/14651858.CD004467.pub2.

- [36] Sahyoun NR, Anderson AL, Kanaya AM, Koh-Banerjee P, Kritchevsky SB, de Rekeneire N, Tylavsky FA, Schwartz AV, Lee JS, Harris TB: **Dietary glycemic index and load, measures of glucose metabolism, and body fat distribution in older adults.** *Am J Clin Nutr* 2005, **82**:547-552.
- [37] Wolever TMS, Jenkins DJA, Vuksan V, Jenkins AL, Wong GS, Joss RG: **Beneficial effect of low-glycemic index diet in overweight NIDDM subjects.** *Diabetes Care* 1992, **15**:562-564.
- [38] Raatz SK, Torkelson CJ, Redmon JB, Reck KP, Kwong CA, Swanson JE, Liu C, Thomas W, Bantle JP: **Reduced glycemic index and glycemic load diets do not increase the effects of energy restriction on weight loss and insulin sensitivity in obese men and women.** *J Nutr* 2005, **135**:2387-2391.
- [39] Young VR, Puig M, Queiroz E, Scrimshaw NS, Rand WM: **Evaluation of the protein quality of an isolated soy protein in young men: relative nitrogen requirements and effects of methionine supplementation.** *Am J Clin Nutr* 1984, **39**:16-24.
- [40] Baglieri A, Mahe S, Zidi S, Huneau JF, Thuillier F, Marteau P, Tome D: **Gastro-jejunal digestion of soya-bean-milk protein in humans.** *Br J Nutr* 1994, **72**:519-532.
- [41] Mariotti F, Mahe S, Benamouzig R, Luengo C, Dare S, Gaudichon C, Tome D: **Nutritional value of [15N]-soy protein isolate assessed from ileal**

- digestibility and postprandial protein utilization in humans.** *J Nutr* 1999, **129**:1992-1997.
- [42] Bos C, Metges CC, Gaudichon C, Petzke KJ, Pueyo ME, Morens C, Everwand J, Benamouzig R, Tome D. **Postprandial kinetics of dietary amino acids are the main determinant of their metabolism after soy or milk protein ingestions in humans.** *J Nutr* 2003, **133**:1308-1315.
- [43] Bhathena SJ, Velasquez MT: **Beneficial role of dietary phytoestrogens in obesity and diabetes.** *Am J Clin Nutr* 2002, **76**:1191-1201.
- [44] Hasler CM: **The cardiovascular effects of soy products.** *J Cardiovasc Nurs* 2002, **16**:50-63.
- [45] Sarkar FH, Li Y: **The role of isoflavones in cancer chemoprevention.** *Frontiers Biosci* 2004, **9**:2714-2724.
- [46] Zhan S, Ho SC: **Meta-analysis of the effects of soy protein containing isoflavones on the lipid profile.** *Amer J Clin Nutr* 2005, **81**:397-408.
- [47] Merz CNB, Johnson BD, Braunstein GD, Pepine CJ, Reis SE, Paul-Labrador M, Hale G, Sharaf BL, Bittner V, Sopko G, Kelsey SF for the Women's Ischemia Syndrome Evaluation Study Group: **Phytoestrogens and lipoproteins in women.** *J Clin Endocrinol Metabol* 2006, **91**:2209-2213.

- [48] Messina M, Ho S, Alekel DL: **Skeletal benefits of soy isoflavones: a review of the clinical trial and epidemiologic data.** *Curr Opin Clin Nutr Metab Care* 2004, **7**:649-658.
- [49] Howes LG, Howes JB, Knight DC: **Isoflavone therapy for menopausal hot flashes: a systematic review and meta-analysis.** *Maturitas* 2006, **55**:302-211.
- [50] Trock BJ, Hilakivi-Clarke L, Clarke R: **Meta-analysis of soy intake and breast cancer risk.** *J Natl Cancer Inst* 2006, **98**:459-471.
- [51] Yan L, Spitznagel EL: **Meta-analysis of soy food and risk of prostate cancer in men.** *Intl J Cancer* 2005, **117**:667-669.
- [52] Jain M, Verma N: **Impact of incorporating soya fibre and processed soyabean flours on the glycaemic index of parantha.** *Asia Pac J Clin Nutr* 2004, **13(Suppl)**:S97.
- [53] Wolever TMS, Radmard R: **Glycemic index of popular nutrition and energy bars.** *J Amer Dietetic Assoc* 2002, **102**:A78.

## Figures

### **Figure 1 - Blood glucose response curves of low-carbohydrate soy products in Experiment 1.**

The mean blood glucose response curves for the equal-carbohydrate portions of the reference food (glucose) and the two soy-based, low-carbohydrate food products tested in Experiment 1. Data are expressed as the change in blood glucose concentration from the fasting baseline concentration. Bars for each data point represent standard error of the means (SEM).

### **Figure 2 - Plasma insulin response curves of low-carbohydrate soy products in Experiment 1.**

The mean plasma insulin response curves for the equal-carbohydrate portions of the reference food (glucose) and the two soy-based, low-carbohydrate food products tested in Experiment 1. Data are expressed as the change in plasma insulin concentration from the fasting baseline concentration. Bars for each data point represent standard error of the means (SEM).

### **Figure 3 - Blood glucose response curves of soy products in Experiment 2.**

The mean blood glucose response curves for the equal-carbohydrate portions of the reference food (glucose) and the four soy-based food products tested in Experiment 2. Data are expressed as the change in blood glucose concentration from the fasting baseline concentration. Bars for each data point represent standard error of the means (SEM).

**Figure 4 - Plasma insulin response curves of soy products in Experiment 2.**

The mean plasma insulin response curves for the equal-carbohydrate portions of the reference food (glucose) and the four soy-based food products tested in Experiment 2. Data are expressed as the change in plasma insulin concentration from the fasting baseline concentration. Bars for each data point represent standard error of the means (SEM).

**Figure 5 - Glycemic and insulin index values of tested soy products**

The mean ( $\pm$  SEM) glycemic index and insulin index for the reference food (glucose) and the six tested soy-based food products. The dark bars represent the glycemic index values and the light bars represent the insulin index values. For the GI values, columns with different superscripts (a, b) are significantly ( $P < 0.001$ ) different. Columns representing the II values with different superscripts (x, y, z) are significantly different ( $P < 0.001$ ).

## Tables

**Table 1 - Mass and nutrient contents of the test portions of the glucose reference and the two test foods in Experiment 1.**

<b>Test Food</b>	<b>Portion Size (g)<sup>1</sup></b>	<b>Energy (kJ [Cal]) {kj/100 g of product}</b>	<b>Protein (g)<sup>2</sup> {g/100 g of product}</b>	<b>Fat (g) {g/100 g of product}</b>	<b>Available Carbohydrate (g) {g/100 g of product}</b>	<b>Sugars (g) {g/100 g of product}</b>	<b>Fiber (g) {g/100 g of product}</b>
Glucose (Reference Food)	10.0 g glucose 250 g water	160 [38.2]	0.0	0.0	10.0	10.0	0.0
Chocolate Raspberry Zing™ bar	150.0	2205 [526.7] {1,470}	45.0 {30.4}	12.5 {8.4}	10.0 {7.7}	0.0 {0.0}	< 2.0 {1.3}
Chocolate	70.0 g powder	1092 [260.8]	40.0	5.0	10.0	2.0	4.0

Daydream™ shake - sucralose	500.0 g water	{1,551}	{57.1}	{7.1}	{14.3}	{2.9}	{5.7}
-----------------------------------	---------------	---------	--------	-------	--------	-------	-------

<sup>1</sup>The amount of water listed for some products is the amount of water needed to prepare that product. This volume is in addition to the 250 g of water consumed with every product as part of the test methodology.

<sup>2</sup>All of the protein in the products above is derived from soy. Each Chocolate Raspberry Zing™ bar contained 21 g soy protein. Each Chocolate Daydream™ shake contained 20 g soy protein.

**Table 2 - Mass and nutrient contents of the test portions of the glucose reference and the four test foods in Experiment 2.**

<b>Test Food</b>	<b>Portion Size (g)</b>	<b>Energy (kJ [Cal]) {kj/100 g of product}</b>	<b>Protein (g)* {g/100 g of product}</b>	<b>Fat (g) {g/100 g of product}</b>	<b>Available Carbohydrate (g) {g/100 g of product}</b>	<b>Sugars (g) {g/100 g of product}</b>	<b>Fiber (g) {g/100 g of product}</b>
Glucose (Reference Food)	25.0 g glucose 250 g water	400 [95.5]	0.0	0.0	25.0	25.0	0.0
Peanut Butter Chocolate Pal™ bar	48.4	813 [194.2] {1,671}	12.9 {26.7}	4.8 {10.0}	25.0 {51.7}	14.5 {30.0}	0.8 {1.7}
Chocolate	47.1 g powder	742	14.7	1.8	25.0	23.6	1.5

Daydream™ shake – fructose	305.0 g water	[177.2] {1,567}	{31.3}	{3.9}	{53.0}	{50.0}	{3.1}
Lightly Salted Sunshine™ soy protein chips	48.1	808 [193/0] {1,672}	13.5 {28.0}	3.8 {8.0}	25.0 {52.0}	0.0 {0.0}	0.0 {0.0}
Soy spaghetti	42.4 (dry)	636 [151.9] {1,467}	10.6 {24.6}	1.1 {2.6}	25.0 {57.9}	1.5 {3.5}	0.8 {1.8}

<sup>1</sup>The amount of water listed for some products is the amount of water needed to prepare that product. This volume is in addition to the 250 g of water consumed with every product as part of the test methodology.

<sup>2</sup>The protein in all of the products except the spaghetti is from soy. For the spaghetti, one serving provided 14 grams of protein, 8 grams from soy and 6 grams from semolina. Each Chocolate Daydream™ shake contained 20 g soy protein. Each Peanut Butter Chocolate Pal™ bar contained 16 g soy protein. The soy protein chips contained 7 g soy protein per serving

**Table 3 - Glycemic index and glycemic load<sup>1</sup> values for the six tested soy food products.**

<b>Test Food</b>	<b>Carbohydrate Tested (g)</b>	<b>Value ± SEM</b>	<b>Category<sup>2</sup></b>	<b>Carbohydrates/ Serving (g)</b>	<b>Value</b>	<b>Category<sup>3</sup></b>
Chocolate Daydream™ shake – sucralose	10.0	25.00 ± 4.28	Low	5	1.25	Low
Chocolate Daydream™ shake – fructose	25.0	32.73 ± 4.41	Low	34	11.13	Medium
Soy spaghetti	25.0	47.03 ± 7.48	Low	33	15.52	Medium
Chocolate Raspberry Zing™ bar	10.0	47.42 ± 4.55	Low	6	2.85	Low
Peanut Butter Pal™ bar	25.0	51.82 ± 3.86	Low	31	16.06	Medium
Lightly Salted Sunshine™ soy pasta chips	25.0	86.79 ± 7.86	High	13	11.28	Medium

<sup>1</sup>Glycemic Load = (GI x net carbohydrates)/100

<sup>2</sup>SEM = Standard Error of the Means

<sup>3</sup>Glycemic Index Category: Low = ≤ 55; Medium = 56 – 69; High = > 70 (Sydney University Glycemic Index Research Service [SUGiRS])

<sup>4</sup>Glycemic Load Category: Low = ≤ 10; Medium = 11 – 19; High = > 20 (SUGiRS; [www.glycemicindex.com](http://www.glycemicindex.com))






Figure 1




Figure 2




Figure 3




Figure 4




Figure 5