

Milk consumption and the prepubertal somatotrophic axis

Janet W. Rich-Edwards^{1,2,3} *
Davaasambuu Ganmaa^{5*}
Michael N. Pollak⁶
Erika K. Nakamoto²
Ken Kleinman²
Uush Tserendolgor⁷
Walter C. Willett^{3,4,5}
Lindsay Frazier^{4,8§}

¹Division of Women's Health, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School

²Department of Ambulatory Care and Prevention, Harvard Medical School and Harvard Pilgrim Health Care

³Department of Epidemiology, Harvard School of Public Health

⁴Channing Laboratory, Brigham and Women's Hospital, Harvard Medical School

⁵Department of Nutrition, Harvard School of Public Health

⁶Division of Cancer Prevention, Department of Oncology, McGill University, Montreal

⁷Public Health Institute, Mongolia

⁸Department of Pediatric Oncology, Dana-Farber Cancer Institute, Harvard Medical School Boston, MA 02115

*Co-first authors

§Corresponding author

Email addresses:

JWRE: jrichedwards@partners.org

DG: gdavaasa@hsph.harvard.edu

MNP: michael.pollak@mcgill.ca

EKN: Erika_Nakamoto@harvardpilgrim.org

KK: ken.kleinman@gmail.com

UT: utseren@mongol.net

WCW: wwillett@hsph.harvard.edu

LF: Lindsay_frazier@dfci.harvard.org

Abstract

Background: Nutrients, hormones and growth factors in dairy foods may stimulate growth hormone (GH), insulin-like growth factor I (IGF-I), and raise the ratio of IGF-I to its binding protein, IGFBP-3. We conducted pilot studies in Mongolia and Massachusetts to test the extent to which milk intake raised somatotrophic hormone concentrations in prepubertal children.

Methods: In Ulaanbaatar, we compared plasma levels before and after introducing 710 ml daily whole milk for a month among 46 10-11 year old schoolchildren. In a randomized cross-over study in Boston, we compared plasma hormone levels of 28 6-8 year old girls after one week of drinking 710 ml lowfat (2%) milk with their hormone levels after one week of consuming a macronutrient substitute for milk.

Results: After a month of drinking whole milk, Mongolian children had higher mean plasma levels of IGF-I ($p < 0.0001$), IGF-I/IGFBP-3 ($p < 0.0001$), and 75th percentile of GH levels ($p = 0.005$). After a week of drinking lowfat milk, Boston girls had small increases in IGF-I, IGF-I/IGFBP-3 and GH, which did not reach statistical significance.

Conclusions: Milk drinking caused increases in somatotrophic hormone levels of prepubertal girls and boys at very different study sites. The finding that milk intake raises GH levels is novel, and suggests that nutrients or bioactive factors in milk may stimulate endogenous GH production. Although the statistically significant Mongolian findings were not macronutrient-controlled, and the macronutrient-controlled Boston findings were not statistically significant, somatotrophic hormone levels were higher among milk drinkers in both studies.

Background

Milk is a complex, bioactive substance honed by evolution to promote growth and development of the infant mammal. Cow's milk and dairy products derived from milk are widely consumed by human children and adults well after the age of weaning. Health benefits of the macronutrient and micronutrient content of milk are established for child development.^[1-6] However, the impact of other bioactive substances in milk has been little investigated. Dairy products contain bovine hormones and growth factors that may alter the hormone and growth factor levels of prepubertal children.

Many bovine enzymes, hormones, and growth factors are selectively compartmentalized in milk.^[10,11] In theory, most protein and peptide hormone and growth factors would be destroyed or deactivated during digestion and first-pass hepatic metabolism. However, several lines of evidence indicate that some hormones or growth factors in milk may survive digestion and circulate in the plasma of milk-drinkers.

Rats fed lowfat (1%) cow's milk from the Japanese grocer had higher levels of IGF-I and estrone sulfate than animals fed a macronutrient substitute for milk.^[16] At least one study of radioactively labeled IGF-I in milk has demonstrated its intact absorption in the rat model.^[17-18] Human studies have consistently shown that high milk consumption is associated with a 10%-20% increase in circulating IGF-I levels among adults and a 20%-30% increase among children.^[3,19-26] Although the protein in milk could be responsible for the higher IGF-I levels in milk-drinkers, dairy products raise IGF-I levels more than meat and other sources of dietary protein.^[20-22, 26] The ratio of IGF-I to its main binding protein, IGFBP-3, also rises with milk consumption, indicating an increase in bioavailable IGF-I.^[19,20] It has recently been hypothesized that dairy consumption may raise IGF-I levels and increase rates of twinning.^[27] Two studies have recently reported that milk intake during pregnancy results in higher birthweight offspring.^[28,29]

Most of the steroid hormones and growth factors produced by cows are identical to those produced by humans.^[30] If bovine hormones survive the digestive process, most are presumably bioactive. To our knowledge, there has been no macronutrient-controlled experimental investigation of the impact of hormones and growth factors in cow's milk on the plasma hormone and growth factor levels of children. We report here

the results of two short-term pilot studies conducted to justify a long-term study of the impact of milk hormones and growth factors on child development.

Methods

1. Mongolian Pilot Study

As children in the US consume large amounts of dairy products, it is difficult to identify a low-dairy intake control group for long-term study of the impact of dairy products on pediatric hormones and development. We turned to Mongolia as a possible site for a long-term study. Mongolia has a long pastoralist tradition, and milk is a staple of herders living on the steppe. A sharp rise in urban migration has resulted in a large group of children whose families lack access to familial herds. Until very recently, the only milk available in the capital city, Ulaanbaatar, was prohibitively expensive imported milk. Ulaanbaatar's children lack access to milk. Lactose intolerance is virtually unknown. In Mongolia, we were able to add milk to children's diets, rather than trying to subtract it from the diets of American children. We report here the results of a study conducted to determine the feasibility of a long-term trial of the impact of milk consumption on child development in Ulaanbaatar.

Participants

With the approval of the Mongolian Ministry of Education and Ministry of Health Ethical Review Board, we conducted a one-month pilot study in Ulaanbaatar from May to June, 2005. The aim of the study was to determine whether participating in research was acceptable to Mongolian parents, children, and institutions, and to test the logistics of shipping milk and blood samples internationally. A letter was sent to the families of all children in a 3rd grade classroom in public school #65 in Ulaanbaatar, which included fifty 9 -11 year old girls and boys. Families were invited to an informational meeting at the school. The Mongolian investigators (Drs. G and T) explained the study protocol and obtained informed consent from parents and assent from children to participate. Children were eligible if they had no known allergy to milk. One child elected not to enroll, two children responded too late to be enrolled, and one child was withdrawn from the study after developing symptoms of milk allergy. A total of 22 girls and 24 boys completed the protocol.

Procedures

We performed a physical examination of each child at the beginning and end of the month, with a Tanner Stage evaluation at the first examination only. At baseline, children completed a food frequency questionnaire regarding their consumption of dairy products in the last seven days. During the intervention month, children consumed three 8-ounce tetrapak “boxes” (710 ml) of conventional U.S. UHT-pasteurized vitamin D fortified whole milk daily. The producer of the milk accepts milk only from dairies that do not use bovine somatotropin (BST). From days 4-10, four children who experienced digestive upset drank varying amounts of milk (236, 472, or 710 ml) as they grew accustomed to the milk. After day 10, all 46 children drank the full 710 ml daily.

Hormone comparisons

At the beginning and end of the month, a morning 10 ml blood sample was drawn into a heparinized tube. Blood samples were immediately chilled, and transported cold to a central laboratory, where serum was separated and stored at -70°C . Samples remained frozen during international shipment. Serum levels of growth hormone, IGF-I, and IGFBP-3 were determined in all samples. We compared pre-intervention and post-intervention hormone levels for each child.

2. Boston Girls and Milk Study

Participants

We conducted a randomized crossover milk feeding study among prepubertal girls, comparing their hormone and growth factor levels after a week of consuming conventional lowfat (2%) milk compared with a week of consuming a vegetable-based macronutrient substitute for milk. The study was conducted at the Boston Children’s Hospital General Clinical Research Center, and was approved by the Children’s Hospital Committee on Clinical Investigation. Thirty-one 6-8 year old girls were recruited from several pediatric clinics and by contacting women who had visited the Cancer Risk and Prevention Clinic of the Dana Farber Cancer Institute. Girls were eligible for the study if their height, weight and body mass index were between the 5th and 95th percentile for age, they had no history of endocrine disorders or clinical signs of pubertal initiation, and they had no allergy to milk or the milk substitute. We obtained consent of parents and assent of the girls for a screening physical and taste-test of the milk substitute. A pediatrician (ALF) performed a brief physical to

ensure that each girl was in Tanner Stage 1, and the girls tested the palatability of the milk substitute before completing a second consent and assent procedure to enroll in the trial. Three girls did not complete the protocol, due to distaste for the milk substitute, fear of blood draw, or inaccessible veins. We present results for 28 girls who completed the trial. Most girls were Caucasian; 2 were African American, 1 was Native American, and 6 were of Asian descent.

Procedures

The protocol was five weeks long, including two intervention weeks and an intervening three-week ‘wash-out’ return to normal diet. In one intervention week, girls consumed 710 ml of 2% conventional cow’s milk per day (from a manufacturer that does not accept milk from cows treated with BST). In a second intervention week, the girls drank 710 ml daily of a macronutrient substitute for milk with the same calorie, protein, fat, carbohydrate, calcium and vitamin D content as 2% cow’s milk, concocted from almond milk, coconut milk, and protein powder. During these two intervention weeks, girls consumed otherwise dairy-free diets. The order of the intervention weeks was randomized. At the introductory visit, parents completed a food frequency questionnaire about their consumption of dairy products over the past year. Girls consumed 99% of the milk and 92% of the milk substitute.

Hormone comparisons

An afternoon 10 ml blood sample was drawn into a heparinized tube at the end of each intervention week, centrifuged, aliquotted, and stored at -70°C . We compared the hormone and growth factor level of each girl at the end of the milk week to her level at the end of the milk substitute week.

Hormone and growth factor analysis for both studies

GH, IGF-I, and IGFBP-3 were assayed by enzyme-linked immunoabsorbent assay (ELISA) with reagents from Diagnostic Systems Laboratory (Webster, Texas). The IGF-I values obtained by ELISA in our laboratory are highly correlated (Pearson $r = 0.97$) with values obtained by radioimmunoassay after acid chromatography. The average intra-assay coefficients of variation for IGF-I, IGFBP-3, and GH were 4.9%, 9.0%, and 8.0%, respectively. The IGF-I assay was run in two batches, with limits of detection of 14 ng/ml and 21 ng/ml, respectively (both samples for each child were included in the same batch). The limits of detection were 2 ng/ml

for the IGFBP-3 assay and 0.2 ng/ml for the GH assay. Results below the limit of detection were assigned the limit of detection for analysis.

Statistical analysis for both studies

We performed paired t-tests of differences in the means of IGF-I and the IGF-I/IGFBP3 ratio; the GH distribution was too skewed to apply a parametric test. We used unpaired quantile regression to test differences in the 50th and 75th percentiles of the distributions. We report the 75th percentile because more than half the GH values were at or below the limit of detection, and the 75th percentile captured variation above this “floor” limit. Paired statistical techniques for quantile regression are not available; however, unpaired analyses are valid but would be less powerful than paired techniques if they existed. To display results graphically, we estimated kernel density distributions of hormone and growth factor levels.^[31] All analyses were performed using SAS (version 9.1) software.

Results

Characteristics of children in the two pilot studies are presented in Table 1. Mongolian children are somewhat smaller and leaner than their U.S. peers, with a profile similar to that of U.S. 9-10 year olds.^[32] The height, weight and body mass index (BMI, kg/m²) of the Boston girls was typical for U.S. girls age 6-8 years.^[32] The Mongolian children consumed an average 0.6 servings of whole milk, 0.7 servings of cheese/week, and 1.6 servings of yoghurt in the week before the intervention. Fifteen percent of children had no dairy intake at all, confirming anecdotal reports of low dairy intake among Mongolian urban children. The Boston girls typically consumed 15 glasses of predominantly lowfat or skim milk per week, with an average 19 servings of other dairy products per week.

Figure 1 shows the kernel density estimated distributions of GH, IGF-I, and IGF-I/IGFBP-3 according to milk intervention arm in each study. As expected, the distribution of GH in both studies was skewed, with the median values in both studies at or near the limit of detection. The highest GH values were observed after milk consumption in both studies. In Mongolia, large increases in IGF-I and IGF-I/IGFBP-3 levels were observed between the beginning and end of the milk intervention month. In Boston, there was a subtle shift of the IGF-I

and IGF-I/IGFBP-3 distributions to higher levels after a week of consuming milk, compared with a week of consuming milk substitute.

Mongolian study

During the milk intervention month, girls grew a mean 1.1 (0.2 sd) centimeters, 0.65 (0.2) kg, and had a mean drop in BMI of -0.29 (0.06). Boys grew a mean 1.0 (0.2) centimeters, 0.23 (0.1) kg, and had a mean decrease in BMI of -0.26 (0.06). Consistent with the rapid increase in height over the intervention month, we observed large increases in plasma GH, IGF-I, and IGF-I/IGFBP-3 (Table 2). Large differences in the means, medians, and 75th percentile of these somatotrophic hormones were statistically significant, with the exception of median GH, which was nearly doubled but was not statistically significant. Changes in GH and growth factor levels from the beginning to the end of the milk intervention month were similar for girls and boys (data not shown).

Table 3 shows the results of statistical tests of the plasma assays in the Boston trial. There were no statistically significant differences in the hormone and growth factor distributions “on” vs. “off” milk, although the mean and the 75th percentile of IGF-I and IGF-I/IGFBP-3, as well as the 75th percentile of GH, were higher in these 6-8 year old girls when they were drinking milk.

Discussion

In the Boston and Mongolian pilot studies, levels of prepubertal GH, IGF-I, and IGF-I/IGFBP-3 were higher after one week and one month, respectively, of drinking milk. Although the macronutrient-controlled Boston findings were not statistically significant, and the statistically significant Mongolian findings were not macronutrient controlled, similar hormonal patterns were observed at both sites. The rise in somatotrophic hormone and growth factor levels was larger in the Mongolian than in the Boston study, perhaps due to the longer intervention, the fact that the Mongolian children consumed less dairy at baseline, the use of whole versus lowfat milk, the older age of the Mongolian participants, or the baseline nutritional status of the Mongolian children.

We conducted the Mongolian feasibility pilot with limited resources, and therefore did not attempt a macronutrient control. During the intervention month, the Mongolian children experienced rapid linear growth (the equivalent of 12 cm/year) compared to an average height velocity of 5-6 cm/year in U.S. children age 10-11 years.^[33] Although such rapid growth suggests that the children may have been undernourished at the start of the trial, the increase in GH is unlikely to be a refeeding effect of milk nutrients. The classic hormonal profile in undernutrition is low IGF-I, but stable or even elevated GH levels.^[34-37] High GH during fasting may promote the tissue-sparing switch from gluconeogenesis to lipolysis during periods of undernutrition.^[34,35] On refeeding, IGF-I levels increase, but GH levels typically fall.^[35,36] In the Mongolian children, the increase in GH levels after milk consumption was proportionally higher than the increase in either IGF-I or IGF-I/IGFBP-3 levels. This finding is inconsistent with a malnutrition correction effect. The similar hormonal response to milk among the well-nourished Boston girls also argues against a refeeding effect. As GH is released in a pulsatile fashion, and we had only one GH measurement in these pilot studies, we undoubtedly overestimated the variability of GH, reducing our statistical power to detect a GH response to milk. Future studies should measure GH more frequently to capture GH secretory dynamics and better estimate GH levels.

Conclusions

Our two pilot studies suggest an immediate, short-term stimulatory impact of milk on the secretion of GH and bioactive IGF-I. Other studies suggest that the impact of milk on IGF-I levels is maintained, and may grow over time. In China, 10 year old girls randomized to consume whole milk had 9% higher IGF-I levels after one year, and 20% higher levels after two years ($p < 0.05$).^[2] In Britain, 12 year-old girls randomized to drink approximately 300 ml/day of milk had 10% higher IGF-I levels after 6 months of intervention, and 17% higher levels at months 12 and 18 of intervention ($p = 0.02$).^[3] As these trials were not designed to measure an impact of non-nutritive factors on hormone levels, they did not include a macronutrient control group. Therefore, the impact of the calories and protein on IGF-I levels cannot be distinguished from the impact of non-nutritive bioactive substances in milk. Although our findings from the macronutrient-controlled Boston study were not statistically significant, the rise in somatotrophic hormones after just one week of milk consumption is the first

suggestion that non-nutritive factors in milk may raise IGF-I levels. To our knowledge, this is also the first study to report growth hormone response to dairy intake, which was large and statistically significant among the Mongolian children.

These findings beg replication in larger studies with macronutrient and micronutrient control. If consistent findings emerge, it would be important to identify the specific bioactive factor in bovine milk that raises GH and IGF-I levels. Although human and bovine GH are not identical, bovine GH absorbed from milk might cross-react with the human GH assay, and it is conceivable (but as yet untested) that bovine GH might stimulate human IGF-I production. Alternatively, another factor in bovine milk may stimulate endogenous production of human GH, which would, in turn, provoke IGF-I production. Although bovine and human IGF-I are identical, IGF-I absorbed from milk is unlikely to stimulate human GH secretion, as GH levels should drop in response to negative feedback from IGF-I infusion. Future analysis should focus on identifying factors in milk that could stimulate human GH secretion, such as growth hormone releasing hormone, ghrelin, or a substance that stimulates ghrelin release.

The findings of the two pilot studies are similar. Taken together, they suggest a sizeable short-term impact of milk on child growth hormone and growth factor levels that may be due to non-nutrient factors in milk. The macronutrient-controlled data from the Boston pilot study suggest that the effect is related specifically to milk, rather than to changes in energy or protein intake. More study is needed to determine whether such short-term elevations in somatotropic hormones are maintained over time. Hormones or growth factor in milk may affect childhood growth, pubertal development, and risk of chronic disease in adulthood. If so, the risks of dairy hormone exposure will need to be weighed against established and emerging nutritional benefits of vitamin-fortified milk.

Abbreviations

BMI = body mass index (kg/m^2)

BST = bovine somatotropin

GH = growth hormone

IGF-I = insulin-like growth factor I

IGFBP-3 = insulin-like growth factor binding protein 3

Competing interests

The authors declare that they have no competing interests.

Authors' Contributions

JRE, DG, LF, WW, and EN designed the study. DG and UT conducted the Mongolian pilot study. LF, JRE, and EN conducted the Boston study. MP contributed hormone assays and interpretation. KK and EN performed statistical analyses. JRE drafted the manuscript, and all contributors have edited and approved the manuscript.

Acknowledgments

These projects were funded by the Breast Cancer Research Foundation; grant M01-RR02172 from the National Center for Research Resources, National Institutes of Health to the Children's Hospital Boston General Clinical Research Center; a translational acceleration grant to Dr. Pollak from the Canadian Breast Cancer Research Alliance; and a charitable contribution by the Boston office of Deloitte & Touche LLP.

We are grateful to the participants in the Boston Girls and Milk Study, the families and staff of Ulaanbaatar public School #65, Mongolian Ministry of Health, Mongolian Ministry of Education, Mongolian Public Health Institute, Dr. Suvdaa, Dr. Jambalmaa, Dr. David Fraser, Mr. Gary Bradwin and Ms. Nyan Chu, Staff of the Boston Children's Hospital General Clinical Research Center, Dr. Henry Feldman of the Clinical Research Program at Boston Children's Hospital, Dr. Judy Garber and staff of the Dana Farber Cancer Risk and Prevention Clinic, and the staff and pediatricians at Longwood Pediatrics and the Children's Hospital Primary Care Clinic who referred patients to our study.

References

- [1] Du X, Zhu K, Trube A, Zhang Q, Ma G, Hu X et al. **School-milk intervention trial enhances growth and bone mineral accretion in Chinese girls aged 10-12 years in Beijing.** *British Journal of Nutrition* 2004, **91**:1-11.
- [2] Zhu K, Du X, Cowell CT, Greenfield H, Blades B, Dobbins TA et al. **Effects of school milk intervention on cortical bone accretion and indicators relevant to bone metabolism in Chinese girls aged 10-12 y in Beijing.** *Am J Clin Nutr* 2005, **81**:1168-1175.
- [3] Cadogan J, Eastell R, Jones N, Barker ME. **Milk intake and bone mineral acquisition in adolescent girls: randomised, controlled intervention trial.** *BMJ* 1997, **315**:1255-1260.
- [4] Kalkwarf HJ, Khoury JC, Lanphear BP. **Milk intake during childhood and adolescence, adult bone density, and osteoporotic fractures in US women.** *Am J Clin Nutr* 2003, **77**:257-265.
- [5] Nicklas TA. **Calcium intake trends and health consequences from childhood through adulthood.** *J Am Coll Nutr* 2003, **22**:340-356.
- [6] Elwood PC, Pickering JE, Hughes J, Fehily AM, Ness AR. **Milk drinking, ischaemic heart disease and ischaemic stroke II. Evidence from cohort studies.** *Eur J Clin Nutr* 2004, **58**:718-724.
- [7] Kuh D, Ben-Shlomo Y. *A Life Course Approach to Chronic Disease Epidemiology*. Oxford: Oxford University Press, 1997.
- [8] Renehan AG, Zwahlen M, Minder C, O'Dwyer ST, Shalet SM, Egger M. **Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis.** *Lancet* 2004, **363**:1346-1353.
- [9] Ganmaa D, Wang PY, Qin LQ, Hoshi K, Sato A. **Is milk responsible for male reproductive disorders?** *Med Hypotheses* 2001, **57**:510-514.
- [10] Grosvenor CE, Picciano MF, Baumrucker CR. Hormones and growth factors in milk. *Endocr Rev* 1993, **14**:710-728.
- [11] Rodriguez-Palmero M, Koletzko B, Kunz C, Jensen R. **Nutritional and biochemical properties of human milk: II. Lipids, micronutrients, and bioactive factors.** *Clin Perinatol* 1999, **26**:335-359.
- [12] Daxenberger A, Breier BH, Sauerwein H. **Increased milk levels of insulin-like growth factor 1 (IGF-1) for the identification of bovine somatotropin (bST) treated cows.** *Analyst* 1998, **123**:2429-2435.
- [13] Epstein SS. **Potential public health hazards of biosynthetic milk hormones.** *Int J Health Serv* 1990, **20**:73-84.
- [14] Epstein SS. **Unlabeled milk from cows treated with biosynthetic growth hormones: a case of regulatory abdication.** *Int J Health Serv* 1996, **26**:173-185.
- [15] Mephram TB, Schofield PN, Zumkeller W, Cotterill AM. **Safety of milk from cows treated with bovine somatotropin.** *Lancet* 1994, **344**:197-198.

- [16] Qin LQ, Xu JY, Wang PY, Ganmaa D, Li J, Wang J, Kaneko T, Hoshi K, Shirai T, Sato A. **Low-Fat Milk Promotes the Development of 7,12-Dimethylbenz(a)anthracene (DMBA)-Induced Mammary Tumors in Rats.** *International Journal of Cancer* 2004, **110**:491-496.
- [17] Philipps AF, Kling PJ, Grille JG, Dvorak B. **Intestinal transport of insulin-like growth factor-I (igf-I) in the suckling rat.** *J Pediatr Gastroenterol Nutr* 2002, **35**:539-544.
- [18] Pacha J. **Development of intestinal transport function in mammals.** *Physiol Rev* 2000, **80**:1633-1667.
- [19] Ma J, Giovannucci E, Pollak M, Chan JM, Gaziano JM, Willett W et al. **Milk intake, circulating levels of insulin-like growth factor-I, and risk of colorectal cancer in men.** *J Natl Cancer Inst* 2001, **93**:1330-1336.
- [20] Giovannucci E, Pollak M, Liu Y, Platz EA, Majeed N, Rimm EB et al. **Nutritional predictors of insulin-like growth factor I and their relationships to cancer in men.** *Cancer Epidemiol Biomarkers Prev* 2003, **12**:84-89.
- [21] Holmes MD, Pollak MN, Willett WC, Hankinson SE. **Dietary correlates of plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 concentrations.** *Cancer Epidemiol Biomarkers Prev* 2002, **11**:852-861.
- [22] Hoppe C, Udam TR, Lauritzen L, Molgaard C, Juul A, Michaelsen KF. **Animal protein intake, serum insulin-like growth factor I, and growth in healthy 2.5-y-old Danish children.** *Am J Clin Nutr* 2004, **80**:447-452.
- [23] Hoppe C, Molgaard C, Thomsen BL, Juul A, Michaelsen KF. **Protein intake at 9 mo of age is associated with body size but not with body fat in 10-y-old Danish children.** *Am J Clin Nutr* 2004, **79**:494-501.
- [24] Rogers IS, Gunnell D, Emmett PM, Glynn LR, Dunger DB, Holly JM. **Cross-sectional associations of diet and insulin-like growth factor levels in 7- to 8-year-old children.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:204-212.
- [25] Heaney RP, McCarron DA, Dawson-Hughes B, Oparil S, Berga SL, Stern JS et al. **Dietary changes favorably affect bone remodeling in older adults.** *J Am Diet Assoc* 1999, **99**:1228-1233.
- [26] Hoppe C, Molgaard C, Juul A, Michaelsen KF. High intakes of skimmed milk, but not meat, increase serum IGF-1 and IGFBP-3 in eight-year-old boys. *Eur J Clin Nutr* 2004;**58**:1211-16.
- [27] Steinman G. **Mechanisms of twinning. VII. Effect of diet and heredity on the human twinning rate.** *J Reprod Med* 2006, **51**:405-410.
- [28] Mannion CA, Gray-Donald K, Koski KG. **Association of low intake of milk and vitamin D during pregnancy with decreased birth weight.** *Canadian Medical Association Journal* 2006, **174**:1273-1277.
- [29] Olsen S, Halldorsson ThI, Willett W, Knudsen VK, Mikkelsen TB, Gillman MW et al. **Association of milk consumption in pregnancy with increased foetal growth.** *Pediatric Research* 2005, **58**:1120.
- [30] Goffin V, Shiverick KT, Kelly PA, Martial JA. **Sequence-function relationships within the expanding family of prolactin, growth hormone, placental lactogen, and related proteins in mammals.** *Endocr Rev* 1996, **17**:385-410.

- [31] Scott DW. *Multivariate Density Estimation: Theory, Practice, and Visualization*. New York: John Wiley & Sons, Inc., 1992.
- [32] Centers for Disease Control and Prevention. **CDC Growth Charts: United States 2000**. Retrieved June 20, 2006 from [<http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm>].
- [33] Tanner JM. *Foetus into Man: Physical Growth from Conception to Maturity*. Cambridge, MA: Harvard University Press, 1990.
- [34] Ho K, Veldhuis JD, Johnson M, Furlanetto R, Evans WS, Alberti KG et al. **Fasting enhances growth hormone secretion in man**. *J Clin Invest* 1988, **81**:968-9753
- [35] Soliman AT, ElZalabany MM, Salama M, Ansari BM. **Serum leptin concentrations during severe protein-energy malnutrition: correlation with growth parameters and endocrine function**. *Metabolism* 2000, **49**:819-825.
- [36] Soliman AT, Hassan AE, Aref MK, Hintz RL, Rosenfeld RG, Rogol AD. **Serum insulin-like growth factors I and II concentrations and growth hormone and insulin responses to arginine infusion in children with protein-energy malnutrition before and after nutritional rehabilitation**. *Pediatr Res* 1986, **20**:1122-1130.
- [37] Hintz RL, Suskind R, Amatayakul K, Thanangkul O, Olson R. **Plasma somatomedin and growth hormone values in children with protein-calorie malnutrition**. *J Pediatr* 1978, **92**:153-156

Table1. Mean (sd) baseline characteristics of participants in two pilot studies.

	Mongolian boys	Mongolian girls	Boston girls
n	24	22	28
Age	10.4 (0.7)	10.6 (0.6)	7.6 (0.9)
Weight (kg)	29.1 (0.7)	28.8 (1.0)	25.6 (4.6)
Height (cm)	132.5 (1.4)	130.8 (1.3)	123.9 (6.6)
BMI (kg /m ²)	16.5 (0.3)	16.8 (0.3)	16.6 (2.0)
Tanner stage >1	12% (3/24)	4% (1/22)	0% (by design)
In past week:			
Servings of milk	0.8 (1.3)	0.5 (1.0)	14.9 (5.8)
Servings of other dairy ¹	6.5 (6.0)	4.1 (4.0)	18.9 (10.3)

1. “Other dairy” for the Boston girls included cheese, ice cream, yoghurt, and cheese-containing dishes such as pizza or pasta. “Other dairy” for the Mongolia children consisted of cheese, yoghurt, or milk added to tea.

Table 2. Change in 10 year-old Mongolian children's hormone and growth factor concentrations before and after one month of drinking 710 ml of American UHT whole milk daily. For means, p-values are from paired t-tests. For 50th and 75th percentiles, p-values are from unpaired quantile regression.

		Mean	50th percentile	75th percentile
Growth Hormone (ng/ml)	Pre	2.38 ± 3.27	1.23	2.60
	Post	3.42 ± 4.07	2.14	5.28
	p-value	NA ¹	0.23	0.005
IGF-I (ng/ml)	Pre	290.93 ± 93.98	270.70	346.95
	Post	358.34 ± 125.62	333.56	449.29
	p-value	<0.0001	0.04	0.03
IGF-I/IGFBP-3	Pre	0.064 ± 0.016	0.062	0.068
	Post	0.073 ± 0.019	0.072	0.082
	p-value	<0.0001	0.009	0.007

¹T-test comparing mean GH not appropriate, due to non-Normal GH distribution.

Table 3. Comparison of hormone and growth factor concentrations among 6-8 year old Boston girls after 1 week milk consumption versus 1 week macronutrient substitute for milk (n=28). For means, p-values are from paired t-tests. For 50th and 75th percentiles, p-values are from unpaired quantile regression.

		Mean	50th percentile	75th percentile
Growth Hormone (ng/ml)	After substitute	0.89 ± 0.96	0.41	1.52
	After milk	1.25 ± 1.67	0.21	1.62
	p-value	NA ¹	0.63	0.93
IGF-I (ng/ml)	After substitute	257.26 ± 69.12	250.36	294.71
	After milk	270.60 ± 89.46	250.20	308.77
	p-value	0.35	0.99	0.68
IGF-I/IGFBP-3	After substitute	0.057 ± 0.01	0.057	0.064
	After milk	0.059 ± 0.02	0.056	0.068
	p-value	0.28	0.72	0.28

¹T-test comparing mean GH is not appropriate, due to non-Normal GH distribution.

Figures

Figure 1 legend - Distributions of plasma somatotrophic hormones in two pilot studies

The Mongolia study compared concentrations of growth hormone (A), insulin-like growth factor I (IGF-I) (B), and the ratio of IGF-I to its binding protein 3 (IGF-I/IGFBP-3) (C) among 9-11 year old Mongolian boys and girls (n=46) before (red line) and after (blue line) one month of drinking 710 ml whole milk daily. The Boston study included 6-8 year old girls (n=28) whose concentrations of growth hormone (D), IGF-I (E), and the ratio of IGF-I/IGFBP-3 (F) were measured after 1 week consuming 710 ml of lowfat milk daily (blue line) and one week of consuming a vegetable-based macronutrient substitute for milk (red line).

Figure 1)

A. Mongolian Growth Hormone






B. Boston Growth Hormone

C. Mongolian IGF-I

D. Boston IGF-I

E. Mongolian IGF-I/IGFBP-3

F. Boston IGF-I/IGFBP-3

A) Mongolian Growth Hormone**B) Boston Growth Hormone****C) Mongolian IGF-I****D) Boston IGF-I****E) Mongolian IGF-I/IGFBP-3****F) Boston IGF-I/IGFBP-3**