

Consumption of Fruits and Vegetables in relation to the Risk of Developing Acute Coronary Syndromes; the CARDIO2000 case-control study

Demosthenes B Panagiotakos^{1,2}, Christos Pitsavos¹, Peter Kokkinos², [Christina Chrysohoou¹](#), ~~Christina Chrysohoou, MD, PhD, Peter F. Kokkinos¹ MD, PhD, FACSM, Nick Marinakis, MD, Peter Stravopodis, MD, Christodoulos Stefanadis, MD, PhD, FESC, FACC and Pavlos Toutouzas, MD, PhD, FESC, FACC~~

~~Section of Preventive Cardiology, Department of Cardiology, School of Medicine, University of Athens, Greece~~

~~¹Veterans Affairs Medical Center and Cardiology Division, Georgetown University, Washington, DC, USA~~

Word count (text):-

Corresponding author:-

~~Demosthenes B Panagiotakos~~

~~48-50 Chiou Str. Glyfada,~~

~~Attica, 165-61, Greece~~

~~Tel. +301-9603116 & +301-9600719 (Fax)~~

~~d.b.panagiotakos@usa.net~~

ABSTRACT

Background. In this work we aimed to evaluate the hypothesis whether coffee consumption is associated with the risk of developing acute coronary syndromes. **Methods.** During 2000 and 2001, 1804 randomly selected subjects from several regions of Greece were entered into a case control study. Of these, 848 were patients with their first acute coronary heart syndrome event, and 1078 were hospitalised controls with no cardiovascular disease in their medical history. The effect of caffeine consumption on coronary risk was examined under the presence of several potential confounders. **Results.** The analysis showed that 83% of male and 70% of female patients as well as 84% of male and 77% of female controls reported that they consume at least one cup of coffee per day (P value = NS). Moreover, controls reported that they consume lower quantities of coffee as compared with patients (P value = 0.001). The multivariate analysis raises a J-shape association between the risk of acute coronary syndromes and the number of consumed cups per day. In particular, the odds ratios for moderate consumption (1–2 cups/day), heavy consumption (3–4 cups/day), and very heavy consumption (≥ 4 cups/day) relative to no consumption were 0.71 (95% confidence interval (CI) 0.56–0.90), 1.61 (95% CI 1.15–2.26), and 4.66 (95% CI 2.33–9.34), respectively, after controlling for the presence of hypertension, hypercholesterolemia, diabetes mellitus, family history of premature coronary heart disease, physical activity status, smoking habits, body mass index, alcohol consumption, feeling of depression and education status. **Conclusion.** This study suggests the J-shape association between coffee consumption and the risk of developing acute coronary syndromes. It seems that moderate coffee drinking benefits human health, whilst heavy consumption is associated with increased coronary risk.

INTRODUCTION

It has long been suspected that coffee consumption may have adverse effects on the cardiovascular system. Some studies indicate a positive association with coronary heart disease [1], whilst others suggested no relationship [2, 3]. In order to explore the pathway by which caffeine consumption may affect coronary risk some cross-sectional studies [5, 6, 7] but not all [8, 9, 10] have attributed the observed positive association to elevated blood pressure levels. On the other hand, based on three decades of follow-up the investigators from the John Hopkins Precursors Study reported that coffee drinking appears to play an insignificant role in the development of hypertension [4]. Furthermore, from a prospective study, Gynelberg et al [3] concluded that the association between coffee consumption and the risk of ischemic heart disease is conditioned by known risk factors correlated with the use of coffee, which may partly explain the inconsistencies in the results of previous studies. However, the independent association between coffee consumption in various quantities and the risk of developing coronary heart disease has not been well established, yet.

In order to test the hypothesis whether coffee consumption is associated with the risk of developing coronary heart disease, we conducted a case-control study, in which we enrolled hospitalised cardiac patients, with confirmed acute myocardial infarction or unstable angina for a first time in their life, and matched by sex, and age cardiovascular disease free individuals, from several Greek regions. The previous association was tested under the presence of the potential confounding effect of several lifestyle factors related to ischemic heart disease and coffee consumption, like the conventional cardiovascular risk factors, the presence of psychological stress and depression, as well as educational status (as a measurement of social level).

METHODS

Study population The CARDIO2000 is a multicentre case-control study that explores the association between several demographic, nutritional, psychological, lifestyle and medical risk factors with the risk of developing non-fatal acute coronary syndromes. In order to reduce the unbalanced distribution of several measured or unmeasured confounders, both patients and controls were randomly selected. A sequence of random numbers (1...0) was applied in the hospitals' admission listings. Thus, the coronary patients who assigned the number 1 were included into the study and interviewed (i.e. approximately half the cardiac

patients that visited each cardiology clinic). The same procedure (i.e. admission listings) was applied for the controls, after taking into account the matching criteria. In the case of population controls the random selection was obtained through the municipal rolls.

From January 2000 to August 2001, 956 individuals who had just entered into the selected hospitals with their first symptoms of coronary heart disease (stable angina was excluded from the analysis), 848 (89%) agreed to be enrolled into the study (cases). The inclusion criteria for cardiac cases included:

diagnosis of first acute myocardial infarction (MI). MI was defined by two features: electrocardiographic changes, compatible clinical symptoms, and specific diagnostic enzyme elevations (49% of the patients had MI); or

diagnosis of unstable angina (i.e. one or more angina episodes at rest within the preceding 48 hours) corresponding to class III of the Braunwald classification (51% of the patients had unstable angina).

After the selection of the cardiac patients 1078 of 1300 (83% response) we randomly selected cardiovascular disease free subjects (controls) among those who agreed to be enrolled into the study. The controls were matched to the patients according their age distribution (within classes of ± 3 years), their sex, and the same Greek region.

Stratification According to the population distribution provided by the Hellenic Statistical Services (census 2000), we stratified our sampling into all the Greek regions, in order to include various socio-economic levels and cultural characteristics of the studied population. The enrolled subjects were from, approximately, half of the clinics (7 public and 2 private) of the two biggest metropolitan cities (Athens and Thessalonica), and from almost all (3/4) the major prefectorial hospitals of the other Greek regions (3 from Sterea Ellada, 3 from Thessalia, 2 from Hpeiros, 5 from Macedonia, 2 from Thrace, 5 from Peloponnese, 2 from Crete, 5 from Aegean and 3 from Ionian islands).

Controls were mainly individuals (91% of the total number of controls) who visited the outpatient departments of the same hospital and at the same period with the coronary patients, for routine examinations or minor surgical operations. However, in country hospitals, where the available number of hospitalised controls was not sufficient for the matching procedure, we enrolled into the study a small number of visitors to the same hospitals (3% of the total number of controls) or individuals (6% of the total number of controls) (randomly selected from municipal rolls). The controls were subjects without any clinical

symptoms, signs or suspicion of cardiovascular disease in their medical history, as evaluated by a cardiologist of the study. We used this type of control in order to have more accurate medical information, to eliminate the potential adverse effect of several, unknown, confounders and to increase the likelihood that cases and controls shared the same study base [11]. The number of subjects was determined through power analysis, in order to evaluate a minimum difference of 7% in exposure history between cases and control with statistical power of 0.80 and $P < 0.05$.

Coffee consumption All participants were asked their usual frequency of consumption of coffee over the last year. Based on the distribution of coffee consumption, we categorized usual caffeine consumption as none, up to 14 cups per week or 1-2 cups per day (moderate use), or up to 28 cups per week or 3-4 cups per day (heavy use) and more than 28 cups per week or more than 4 cups per day (very heavy use). All reported types of coffee (instant coffee, «Greek», «French» or «cappuccino») were adjusted for one cup of 150ml coffee and concentration of 27.5% caffeine. We did not determine the consumption of decaffeinate coffee, tea and caffeine containing drinks (coca-cola) or chocolate consumption.

Other exposure parameters We defined as current smokers those who smoked at least one cigarette per day. Former smokers were defined as the subjects who had stopped smoking for over 1 year. Individuals who reported that they had never smoked a cigarette in their life were considered as never smokers. Occasional smoking was recorded, and occasional smokers excluded from the analysis due the small number in cases (12, 1%) and controls (21, 2%). Educational level was measured by years of schooling. Mean annual income, during the past 5 years, was also recorded. Clinical symptoms of depression during the previous month were sought using the Center of Epidemiological Studies Depression scale (CES-D). Based on a pilot analysis in clinically confirmed depressive subjects from the same study base, a cut off point of 15 on the scale (0-60) was used in order to discriminate significant depressive symptoms. Thus, we classified the subjects into two categories, those with and without short term depressive symptoms (CES-D > 15 or < 15) [12]. Physical activity was defined as any type of non-occupational physical exercise, at least once per week during the past year. The rest of the subjects were defined as physically inactive. Also, the duration of physical exercise was taken into account. The evaluation of the nutritional habits was based on a

~~questionnaire from the Department of Nutrition of the National School of Public~~ Manolis Vavouranakis¹,
Christodoulos Stefanadis¹, and Pavlos Toutouzas¹

¹Department of Cardiology, School of Medicine, University of Athens, Greece

²Veterans Affairs Medical Center and Cardiology Division, Georgetown University, Washington, DC, USA

Corresponding author:

Demosthenes B Panagiotakos

46 Paleon Polemiston & Heraklitou St. Glyfada,

Attica, 166 74, Greece

Tel. +30210-9603116 & +30210-9600719 (Fax)

d.b.panagiotakos@usa.net

Abstract

Background. The relation between diet and human health has long been investigated. The aim of this work is to evaluate the association between CHD risk and the consumption of fruit and vegetable, in a large sample of cardiac patients and controls. **Methods.** Stratified sampling from all Greek regions, consisted of 848 (700 males, 58 ± 10 years old and 148 females, 65 ± 9 years old) randomly selected patients, admitted to the cardiology clinic for a first event of an acute coronary syndrome (ACS). In addition we selected 1078 frequency paired, by sex-age-region, controls in the same hospitals but without any clinical suspicion of CHD. Using validated food-frequency questionnaires we assessed total diet, including fruit and vegetable intake, on a weekly basis. Multiple logistic regression analysis estimated the relative risk of developing ACS by level of fruits and vegetables intake after taking into account the effect of several potential confounders. **Results.** Data analysis revealed that the benefit of fruit or vegetable consumption increases proportionally by the number of servings consumed (P for trend < 0.001). After adjusting for the conventional cardiovascular risk factors, those in the upper quintile of fruit consumption (5 or more items/day) had 72% lower risk for CHD (odds ratio = 0.28, 95% CI 0.11 - 0.54, $P < 0.001$), compared with those in the lowest quintile of intake (< 1 items /day). Similarly, consumption of vegetable more than 3 days / week was associated with 70% lower risk for CHD (odds ratio = 0.30, 95% CI 0.22 - 0.40, $P < 0.001$), compared with those that they did not consume vegetables. Of particular interest, a 10% reduction in coronary risk was observed for every one piece of fruit consumed per day (odds ratio = 0.90, 95% CI 0.85 - 0.97, $P = 0.004$). **Conclusions:** Consumption of fruits and vegetables seems to offer significant protection against CHD.

Background

Coronary heart disease is one of the leading causes of death, universally and one of the most common chronic illnesses in many countries of the developing world [1, 2]. Among the conventional cardiovascular risk factors, diet poses a particular challenge for research. Cross-cultural comparisons, case-control and prospective observational studies identified a relationship between diet, blood pressure and lipids levels [3-6], but there is still considerable scientific uncertainty about the relationship between specific dietary components and cardiovascular risk [7], especially in Mediterranean populations.

In this work we quantified the public health benefit of fruit and vegetable consumption on the primary prevention of acute coronary syndromes (acute myocardial infarction or unstable angina) using a large sample of cardiac patients and frequency-matched controls from all Greek regions.

Methods

The CARDIO2000 is a multicentre case-control study that explores the association between several demographic, nutritional, lifestyle, clinical and biochemical risk factors with the risk of developing non-fatal acute coronary syndromes. From January 2000 to March 2002, we randomly selected 848 coronary patients and 1078 cardiovascular symptom free subjects (controls). The number of the enrolled subjects was decided through power analysis, in order to evaluate (two sided) differences in the coronary relative risk greater than 7% with statistical power greater than 80% and significant level less than 5%. According to the population distribution provided by the National Statistical Services (Ministry of Economics, census 1991), we stratified our sampling into all the Greek regions. Controls were mainly individuals who visited the outpatient departments of the same hospital at the same period with the coronary patients, for routine examinations or minor surgical operations. In a few cases (in country hospitals) where the available number of hospitalised controls was not sufficient for the matching procedure, we enrolled into the study friends or colleagues of the coronary patients. All participants were informed about the aims of the study and agreed to give the requested information. In order to eliminate recall bias we tried to retrieve precise information from cases and controls and medical history through hospital or the insurance records.

The coronary patients were selected from the admission listing of the cardiological clinics. According

to the study's protocol the patients were hospitalised for an acute first event of coronary heart disease (acute myocardial infarction or unstable angina; stable angina was excluded from the analysis). The selection criteria for cardiac patients are:

- diagnosis of first acute myocardial infarction (MI). MI was defined by any two features: electrocardiographic changes, compatible clinical symptoms, and specific diagnostic enzyme elevations of SGOT, CPK, LDH and troponin T or I (49% of the patients had MI); or
- diagnosis of unstable angina (i.e. one or more angina episodes at rest within the preceding 48 hours) corresponding to class III of the Braunwald classification (51% of the patients had unstable angina).

Afterwards, we randomly selected subjects without any clinical symptoms, signs or suspicions of cardiovascular disease in their medical history (controls), frequently matched to the patients by age (± 3 years), sex, and region. The CARDIO2000 investigators performed the evaluation of clinical symptoms and signs in the series of cases and controls.

The information regarding the investigated medical factors was retrieved from the subjects' medical records, along with lifestyle characteristics, through a confidential, detailed questionnaire administered during Physician interview after the second day of hospitalisation, for cases and at entry for controls. The subjects social status was investigated through educational and financial stratification. In particular, the educational level was classified into three groups: Group I: education through high school (0 – 9 years of schooling); Group II: high school, technical education up to 14 years of schooling (but not academic) and Group III: (university). Current smokers were defined as those who smoked at least one cigarette per day. Former smokers were those who had stopped smoking for at least one year, and never smokers those who never smoked a cigarette in their lives. Those who stopped smoking for less than a year were classified as smokers. Quantification of smoking status was based on the calculation of pack-years adjusted for nicotine content equal to 0.8 mg per cigarette. Physical activity was defined as any type of non-occupational physical exercise, at least once/week during the past year, and was graded in qualitative terms such as light (expended calories < 4 Kcal/ min, i.e. walking slowly, stationary cycling, light stretching etc.), moderate (expended calories 4-7 Kcal/ min, i.e. walking briskly, outdoor cycling, swimming moderate

effort etc.) and vigorous (expended calories >7 Kcal/ min, i.e. walking briskly uphill, long distance running, cycling fast or racing, swimming fast crawl etc.). The rest of the subjects were defined as physically inactive. Moreover, since physical activity is an important factor for cardiovascular disease, we took into account the occupational activity status of all participants.

The evaluation of the nutritional habits was accomplished with a validated questionnaire based on the guidelines from the Department of Nutrition of the National School of Public Health [8]. We measured the consumption of several food items as an average per week during the past year. The frequency of consumption was quantified approximately in terms of the number of times per month this food was consumed. Thus, daily consumption multiplied by 30 and weekly consumption multiplied by 4 and a value of 0 was assigned to food items rarely or never consumed. In order to describe overall diet we used composite scores, which are necessary for the evaluation of epidemiological associations. Specifically for fruit and vegetable intake the participants reported their average consumption of a specific portion size for these foods over the past year, including apples, grapes, pears, bananas, oranges, etc, as well as broccoli, carrots, spinach, lettuce salad, yellow squash, and tomatoes. Vegetable intake was categorized into <1 serving/day, 1 – 1.49 servings/day, 1.5 – 1.99 servings/day, 2 – 2.49 servings/day, and 2.5+ servings/day, based both on maintaining a natural gradient of exposure and including adequate participants in each category. The provided list of fruits and vegetables was comprehensive enough to capture most types of fruits and vegetables that our subjects consumed. Although fruit juice was recorded it was not evaluated due to low consumption ($< 5\%$). Alcohol, a potential confounder for total energy intake, was measured by daily ethanol consumption (red or white wine, beer, and other beverages), in wineglasses of 100 ml and 12% ethanol concentration) as done by other investigators [9].

According to the collected medical information, the majority of the controls (86%) and the patients (83%) had at least one laboratory measurement during the past 12 months. In addition, we measured, both in patients and controls, arterial blood pressure levels in the right arm (average of 3 measurements having the patient seated and rested), total cholesterol and fasting glucose levels. For the cardiac patients the measurements were collected during the first 12 h of hospitalization and at the end of the interview for the controls. The previous information as well as the patients' and controls' reports assisted us in characterizing the subjects as having hypertension, hypercholesterolemia, or diabetes. In keeping with the long-standing

classification criteria used in several population-based studies, patients whose blood pressure, according to their medical records, were greater or equal to 140 / 90 mmHg or were taking antihypertensive medication were classified as hypertensives. Hypercholesterolemia was defined as cholesterol levels greater than 220 mg/dl or greater than 200mg/dl when two other risk factors for coronary heart disease were present or if hypo-lipidemic treatment was administered. Diabetics were those with fasting blood glucose greater than 125 mg/dl or those who were under special diet or treatment. Finally, we measured the height and the weight both in patients and controls and we calculated the body mass index ($BMI = \text{weight} / \{\text{height}\}^2$). In addition, we asked for any significant changes in their body mass during the past years. Obesity was defined as $BMI > 29.9 \text{ Kg} / \text{m}^2$. Presence of premature coronary heart disease among first-degree relatives (< 55 years for male relatives and < 65 for female relatives) was also recorded. Further details regarding the aims, design and methodology of the CARDIO2000 study have been previously presented [10 - 13].

Statistical analysis. Continuous variables are presented as mean \pm one standard deviation, while qualitative variables are presented as absolute and relative frequencies. In order to fit a multivariate risk model that evaluates the investigated parameters on the risk of developing acute coronary syndromes, an exploratory analysis was initially applied. In particular, stratified by sex contingency tables were conducted in order to investigate associations between categorical variables and groups of study, by the calculation of the overall chi-squared criterion. Also, Wilcoxon non-parametric criterion was applied in order to evaluate differences in continuous measurements (i.e. body mass index) between groups. Estimations of the relative risks of developing acute coronary syndromes were performed by the calculation of the odds ratio (OR) and the corresponding confidence intervals through multiple conditional logistic regression analysis, adjusting for age (in years), body mass index (BMI) (kg/m^2), smoking, alcohol intake, physical activity, history of high cholesterol or hypertension or diabetes or premature CHD, and use of multivitamins. We then conducted a stratified analysis according to smoking status (past, current, never) to evaluate whether the relation between fruits and vegetable intakes and CHD risk differed by these variables. Tests of linear trend across increasing categories of vegetable consumption were conducted by assigning the medians of intakes in categories (servings / day) treated as a continuous variable.

All reported P-values are from two-sided tests and compared to a significant level of 5%. STATA

6 software was used for the calculations (STATA Corp. College Station, Texas, USA).

Results

Seven hundred (83%) of the coronary patients were males and 148 (17%) were females, while 862 (80%) of the controls were males and 216 (20%) were females. Table 1 describes the characteristics of the patients and the controls. The data analysis revealed that 365 (43%) of the cardiac patients and 722 (67%) of the controls consumed some quantity of fruit and vegetable daily. Of these, 28 (3%) patients and 15 (1%) controls reported consuming < 1 serving/day; 351 (41%) patients and 377 (35%) controls reported consuming 1 - 1.49 servings/day; 344 (40%) patients and 428 (40%) controls reported 1.5 – 1.99 servings/day; 100 (9%) patients and 150 (14%) controls reported 2 – 2.49 servings/day, and 33 (7%) patients and 100 (10%) controls consumed 2.5+ servings/day (P for trend < 0.001). We then examined the association between the consumption of fruit, or vegetable and the prevalence of hypertension, diabetes mellitus, hypercholesterolemia, as well as cigarette smoking, body mass index, and physical inactivity, since these factors are considered modifiers of the effect of fruits and vegetables on coronary risk [9]. After stratifying our analysis by group of study (patients vs. controls), an inverse association revealed between fruit and vegetable consumption and cigarette smoking (P-value = 0.008, P-value = 0.04, respectively), prevalence of hypertension (P-value = 0.038, and P = 0.067, respectively), and diabetes mellitus (P = 0.044, and P-value = 0.04, respectively). Furthermore, a positive association was found between fruit and vegetable consumption and education level both in patients and controls (P-value = 0.042, P-value = 0.001, respectively). No significant associations were observed between fruit and vegetables intake and the prevalence of hypercholesterolemia (P = 0.574, P-value = 0.804, respectively), obesity (P-value = 0.579, P-value = 0.565, respectively), alcohol consumption (P-value = 0.939, P-value = 0.919, respectively) and physical inactivity status (P-value = 0.384, P-value = 0.067, respectively). After adjusting for the aforementioned risk factors, those in the upper quintile of fruit consumption (5 or more items /day) had 72% lower relative risk for CHD (OR=0.28, 95% CI 0.11 - 0.54), compared with those in the lowest quintile of intake (<1 item /day). Similarly, consumption of vegetable more than 3 days / week was associated with 70% lower relative risk for CHD (odds ratio = 0.30, 95% CI 0.22 - 0.40), compared with those that they did

not consume vegetables. Of particular interest, a 10% reduction in coronary risk was observed for every additional piece of fruit consumed per day (OR = 0.90, 95% CI 0.85 - 0.97), after controlling for the conventional cardiovascular risk factors. Adjusting for the same covariates in an analysis of the overall trend, considering vegetable intake as a continuous variable, we found an odds ratio of 0.81 (95% CI 0.67 – 0.93) for developing CHD per each additional serving / day of vegetable consumed. The individual effect of fruit and vegetable consumption on coronary risk is presented in Table 2. Moreover, green vegetable intake showed a 12% reduction on coronary risk per item/day (odds ratio = 0.88, 95% CI 0.77 - 0.97), and vitamin C-rich fruits showed an 11% reduction per item/day (odds ratio = 0.89, 95% CI 0.84 - 0.95).

We then stratified the previous analysis by smoking status (current vs. never / former smokers) and found that the protective effect of fruit and vegetable consumption on coronary risk is diminished in the subgroup of heavy (+31 cigarettes per day) current smokers. However, some protection appears to be provided. More specifically, consumption of 2.5+ servings of vegetable per day is associated with a 27% lower risk of developing acute coronary syndromes compared to those who consumed <1 serving per day (odds ratio = 0.73, 95% CI 0.27 – 1.96). Moreover, each 1-piece/day increase of fruit consumption is associated with 5% lower coronary risk in current smokers (odds ratio = 0.95, 95% CI 0.86 – 1.04).

Discussion

The present study revealed that even a moderate consumption of fruit and vegetable (1 – 2 servings per day) is associated with significantly lower risk of coronary events, even after controlling for several potential confounding risk factors (Table 2). Furthermore, the risk was progressively lower as the consumption of fruits or vegetables increased. Those in the highest quintile of consumption (2.5 or more serving per day) had about 70% lower relative risk when compared to those in the lowest quintile. Moreover, some protection appears to be provided even in the group of current smokers.

Our findings are in accordance with those of previous epidemiological studies, which concluded that the consumption of fruits and vegetables might reduce disease risk, particularly for cancer and cardiovascular disease [9, 14, 15]. However, less is known about the direct association between fruits and vegetable intake and risk of acute coronary syndromes [9, 15]. Nine of ten ecological studies two out of

three case-control studies and six of sixteen cohort studies reported a significant protective association of CHD with consumption of fruit and vegetables [5]. Moreover, a review of 250 observational studies reported that increased consumption of fruits and vegetables could reduce cardiovascular deaths from 6% to 22% (8,000 deaths annually) [16]. Our findings support that even low consumption of fruits and vegetables (1-2 servings per week) is associated with about 45% lower coronary risk (Table 2). Consumption of 2 or more servings per week is associated with about 70% reduction in relative risk. This translates to a considerable prevention of acute coronary events, carries a large public health potential, and emphasises the importance of following dietary recommendations as a whole. Moreover, our findings on smokers and the consumption of fruits and vegetables are worth mentioning. Although statistical significance was not achieved, a trend was evident, suggesting that the consumption of fruits and vegetables may offer some protection for current smokers. Since smokers consume substantially less fruit and fruit juice than non-smokers with heavy smokers eating the least [17], special attention should be given to dietary interventions for smokers in order to improve their nutrition. However, smoking cessation must remain the primary target for CHD prevention. A discussion of the mechanisms responsible for the protection fruit and vegetable consumption offers is beyond the scope of this study. However, several investigators reported that the antioxidant, potassium, fiber, and folate content of fruits and vegetables could at least partially explain their protective effect of CHD [18 - 20]. Other compounds such as flavonoids, phytates, lycopene, carotenoids, and other phytochemicals in vegetables may also have significant protective effects in reducing coronary risk [21].

Despite our attempt to control for several known confounding factors, it is still possible that the observed inverse association between fruits and vegetable intake and coronary risk could be at least partially explained by other factors associated with heart-healthy behaviours [22]. Also, in retrospective case-control studies two main sources of systematic errors may exist, the selection and the recall bias. In order to eliminate selection bias we set objective criteria both for patients and controls. However, insignificant misclassification may exist, since a small percentage of asymptomatic coronary patients may be wrongly assigned to controls, even they were evaluated by a cardiologist. Moreover, in case-control studies it is usually observed that patients who had a recent adverse event are more likely to place greater emphasis on several factors related to the disease than the control group (recall bias). To reduce this type of

bias and analyse precise information, we obtained accurate information from the patients as well as from their relatives or their accompanying persons and compare their responses with Kendall's tau-criterion. Concerning the medical information we avoided recall bias by obtaining detailed data from subjects' medical records. However, over/under estimation of the relative risk may exist, especially due to the measurement of nutritional and smoking habits and the onset of the investigated cardiovascular risk factors. Moreover, the coronary patients who died at entry or the day after were not included into the study. This bias could influence our results. However, the proportion of deaths during the first two days was less than 4%. Thus, it is unlikely that the exclusion of fatal events has significantly altered our findings. Finally, we tried to reduce the potential effect of uncontrolled or unknown confounders using the same study base, for patients and controls.

Conclusion

The presented scientific evidences are sufficient to give good reason for public health education and promotion aimed at a substantial increase in the consumption of fruits and vegetables. However, our data do not suggest evidences for causality and population trials are needed in order to provide scientific proof of their efficacy.

Competing interests None declared

Authors' contributions

DP: design of the study, statistical analysis and drafted the manuscript, CP: design of the study and drafted the manuscript, PK: editing, CC: drafted the manuscript, MV: editing, CS: conception of the study and editing, and PT: conception of the study and editing. All authors read and approved the final manuscript.

Acknowledgment

This study is supported by research grants from the Hellenic Heart Foundation (11/1999-2002). The authors would like to thank the physicians and the specialists that coordinated the data collection of this study: Dr K

Tzioumis (Athens, Crete, Peloponisos), Dr J Skoumas (Athens), Dr N Papaioannou (Athens, Thessalia), Dr P Starvopodis (Ionian Islands), Dr L Karra (Aegean Islands), Dr D Antoniadis (Macedonia), Dr G Rembelos (Aegean Islands), Dr D Markou (Athens), A Moraiti (Athens), D Evagelou (Crete), Dr S Vellas (Attica, Hpeirous), Dr. G Skoumbourdis (Stereia Hellas), B Meidanis (Macedonia, Sterea Hellas, Thessalia), Dr S Loggos (Attica), Dr G Zavitsanakis (Stereia Hellas), Dr I Elefsiniotis (Athens), Dr A Theodorakis (Stereia Hellas), Dr N Marinakis (Aegean Islands), Dr T Kyratzoglou (East Macedonia) and S Zombolos (Peloponnese).

References

1. Wilson P, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. **Prediction of Coronary Heart Disease Using Risk Factor Categories.** *Circulation*, 1998; 97:1837-1847.
2. Wood D. **Established and emerging cardiovascular risk factors.** *Am Heart J* 2001; 141:49-57.
3. Keys A, Menotti A, Karvonen MJ, et al. **The diet and 15-year death rate in the seven countries study.** *Am J Epidemiol* 1986; 124:903-915
4. Trichopoulou A, Naska A, Vasilopoulou E. **Guidelines for the intake of vegetables and fruit: the Mediterranean approach.** *Int J Vitam Nutr Res* 2001;71(3):149-53.
5. Ness AR, Powles JW. **Fruit and vegetables, and cardiovascular disease: a review.** *Int J Epidemiol* 1997; 26:1-13.
6. Law MR, Morris JK. **By how much does fruit and vegetable consumption reduce the risk of ischaemic heart disease?** *Eur J Clin Nutr* 1998;52:549-56.
7. Boushey C J, Beresford S A A, Omenn G S, Motulsky A G. **A quantitative assessment of plasma homocysteine as a risk factor for vascular disease: probable benefits of increasing folic acid intakes.** *JAMA* 1995; 274: 1049-57.
8. Trichopoulou A. **From Research to Education: The Greek experience.** *Nutrition* 2000; 16: 528 - 531.
9. Trichopoulou A, Lagiou P, Trichopoulos D. **Traditional Greek diet and coronary heart disease.** *J Cardiovasc Risk* 1994; 1: 9-15.
10. Panagiotakos DB, Pitsavos C, Chrysohoou C, Ioanna Papaioannou, Stefanadis C, Toutouzas PK. **Risk Stratification of Coronary Heart Disease, In Greece: Final Results from CARDIO2000 Epidemiological Study.** *Prev Med* 2002; 35; 548-556.
11. Panagiotakos DB, Pitsavos C, Chrysohoou C, Tzioumis K, Papaioannou I, Stefanadis C, Toutouzas PK. **The Association Of Mediterranean Diet With Lower Risk Of Acute Coronary Syndromes, In Hypertensive Subjects.** *Int J Cardiol* 2002; 82; 141-147.
12. Pitsavos C, Panagiotakos DB, Chrysohoou C, Skoumas J, Papaioannou I, Stefanadis C, Toutouzas PK. **The Benefits From Mediterranean Diet On The Risk Of Developing Acute Coronary**

- Syndromes, In Hypercholesterolemic Subjects: A Case-Control Study (Cardio2000).** *Cor Artery Dis* 2002; 13: 295 – 300.
13. Chrysohoou C, Panagiotakos DB, Pitsavos C, Kokkinos P, Marinakis N, Ioanna-Papaiannou Stefanadis C, Toutouzas PK. **Gender Differences on the Risk Evaluation of Acute Coronary Syndromes: The CARDIO2000 Study.** *Prev Cardiol* 2002;
 14. Willett W. **Nutritional Epidemiology.** *New York: Oxford. University Press*, 1990.
 15. Marchand JL, Luce D, Goldberg P, Bugel I, Salomon C, Goldberg M. **Dietary factors and the risk of lung cancer in New Caledonia (South Pacific).** *Nutr Cancer* 2002; 42(1):18-24.
 16. Van't Veer P, Jansen MC, Klerk M, Kok FJ. **Fruits and vegetables in the prevention of cancer and cardiovascular disease.** *Public Health Nutr* 2000; 3(1):103-7.
 17. Birkett NJ. **Intake of fruits and vegetables in smokers.** *Public Health Nutr* 1999; 2(2): 217-22.
 18. Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC. **Dietary antioxidant intake and risk of coronary heart disease among men.** *N Engl J Med* 1993; 328:1450–56.
 19. American Heart Association. **Dietary Guidelines for healthy American Adults.** *Circulation* 1996; 94:1795–800.
 20. Knekt P, Jarvinen R, Reunanen A, Maatela J. **Flavonoid intake and coronary mortality in Finland: a cohort study.** *Br Med J* 1996;312: 478–81.
 21. Liua S, Leea IM, Ajania U, Colea SR, Buring JE, Mansona JA. **Intake of vegetables rich in carotenoids and risk of coronary heart disease in men: The Physicians' Health Study.** *Int J Epidemiol* 2001;30:130-135.
 22. Nestle M. **Fruits and vegetables: protective or just fellow travelers?** *Nutr Rev* 1996; 54(8): 255-7.

Table 1. Characteristics of the study's population

	ACS Patients				Controls				P-value
	Males		Females		Males		Females		
Number	700 (82%)		148 (18%)		862 (80%)		216 (20%)		
Age (years)	59.1 ± 10		65.3 ± 9		58.8 ± 10		64.8 ± 10		
Education									0.012
<i>Group I (0 – 9 yrs)</i>	406	58%	114	77%	474	55%	147	68%	
<i>Group II (10 – 14 yrs)</i>	182	26%	31	21%	198	23%	41	19%	
<i>Group III (> 14 yrs)</i>	112	16%	3	2%	190	22%	28	13%	
Current smoking	525	75%	44	30%	500	58%	54	25%	< 0.001
Pack years	39.9 ± 14		13.1 ± 9		21.5 ± 13		5.2 ± 6		< 0.001
Hypertension	308	44%	102	69%	216	25%	69	32%	< 0.001
Hypercholesterolemia	413	59%	40	27%	586	68%	67	31%	< 0.001
Diabetes mellitus	168	24%	15	10%	267	31%	17	8%	< 0.001
Family history of CHD	308	44%	77	52%	129	15%	39	18%	< 0.001
Body mass index (Kg/m ²)	27.4 ± 4		27.1 ± 4		27.1 ± 3		26.7 ± 2		0.120
Physical activity	252	36%	37	25%	371	43%	84	39%	0.003
Alcohol consumption	336	48%	25	17%	379	44%	30	14%	0.024

Comparisons were based on the chi-squared test and the Wilcoxon criterion. P-value is for group differences after controlling for gender.

Table 2. Results from the multivariate logistic model that was developed for the evaluation of the effect of fruits and vegetables consumption on coronary risk

Variable	OR	95% C.I. for OR		P-Value
		Lower	Upper	
Age (years)	1.015	1.006	1.025	0.001
Female gender	0.402	0.320	0.505	< 0.001
Vegetable consumption				
<1 serving / day	1.00	-	-	-
1 – 1.49 servings / day	0.562	0.387	0.901	0.012
1.5 – 1.99 servings / day	0.367	0.189	0.712	0.003
2 – 2.49 servings / day	0.297	0.150	0.588	< 0.001
2.5+ servings / day	0.281	0.110	0.541	< 0.001
Fruit consumption (per item)	0.904	0.845	0.967	0.004

The odds ratios were estimated after taking into account the effect of hypertension, hypercholesterolemia, diabetes mellitus, family history of premature coronary heart disease, physical activity status, smoking habits, body mass index, alcohol consumption, and education status of the participants.