

**Development of estimates of dietary nitrates, nitrites, and nitrosamines for use with the Short Willet Food Frequency Questionnaire**

John S. Griesenbeck<sup>1</sup>, MPH  
Michelle D. Steck<sup>2</sup>, BS  
John C. Huber Jr.<sup>3</sup>, PhD  
Joseph R. Sharkey<sup>4</sup>, PhD, RD  
Antonio A. Rene<sup>5</sup>, PhD  
Jean D. Brender<sup>6\*</sup>, PhD

<sup>1</sup>Department of Epidemiology and Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. jsgriesenbeck@srph.tamhsc.edu. Phone: 979-845-2675, fax: 979-458-1877

<sup>2</sup>Department of Epidemiology and Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. mdsteck@srph.tamhsc.edu. Phone: 979-845-2675, fax: 979-458-1877

<sup>3</sup>Department of Epidemiology and Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. jchuber@srph.tamhsc.edu. Phone: 979-862-3432, fax: 979-458-1877

<sup>4</sup>Department of Social and Behavioral Health, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. jrsharkey@srph.tamhsc.edu. Phone: 979 458-4268, fax: 979 458-4264

<sup>5</sup>Department of Epidemiology and Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. aarene@srph.tamhsc.edu. Phone: 979-458-2239, fax: 979-458-1878

<sup>6</sup>Department of Epidemiology and Biostatistics, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. jdbrender@srph.tamhsc.edu. Phone: 979-862-1573, fax: 979-458-1877

\*Corresponding Author: Joseph R. Sharkey, Department of Social and Behavioral Health, School of Rural Public Health, Texas A&M Health Science Center, College Station, TX 77843-1266. jrsharkey@srph.tamhsc.edu. Phone: 979 458-4268, fax: 979 458-4264

The authors declare no conflicts of interest.

Word count:

Title page: 254

Abstract: 255

Text and references: 4,447

Number of Figures: 0

Number of Tables: 3

Running Title: Nitrates, nitrites, & nitrosamines in foods

## **Abstract**

**Background:** Studies have suggested that nitrates, nitrites, and nitrosamines have an etiologic role in adverse pregnancy outcomes and chronic diseases such as cancer. Although an extensive body of literature exists on estimates of these compounds in foods, the extant data varies in quality, quantified estimates, and relevance. **Methods:** We developed estimates of nitrates, nitrites, and nitrosamines for food items listed in the Short Willet Food Frequency Questionnaire (WFFQ) as adapted for use in the National Birth Defects Prevention Study. Multiple reference databases were searched for published literature reflecting nitrate, nitrite, and nitrosamine values in foods. Relevant published literature was reviewed; only publications reporting results for items listed on the WFFQ were selected for inclusion. The references selected were prioritized according to relevance to the U.S. population. **Results:** Based on our estimates, vegetable products contain the highest levels of nitrate, contributing as much as 189 mg/serving. Meat and bean products contain the highest levels of nitrites with values up to 1.84 mg/serving. Alcohol, meat and dairy products contain the highest values of nitrosamines with a maximum value of 0.531  $\mu$ g/serving. The estimates of dietary nitrates, nitrites, and nitrosamines generated in this study are based on the published values currently available. **Conclusions:** To our knowledge, these are the only estimates specifically designed for use with the adapted WFFQ and generated to represent food items available to the U.S. population. The estimates provided may be useful in other research studies, specifically in those exploring the relation between exposure to these compounds in foods and adverse health outcomes.

## **Introduction**

Studies have suggested that nitrates, nitrites, and nitrosamines have an etiologic role in adverse pregnancy outcomes and other health conditions [1-5]. Nitrates and nitrites are precursors in the formation of *N*-nitroso compounds, a class of genotoxic compounds consisting of nitrosamines and nitrosamides [2]. *N*-nitroso compounds are known to cause congenital malformations in animal models, and the role of these compounds in adverse pregnancy outcomes warrants further exploration [6-8]. Humans are exposed to nitrates primarily through diet and drinking water, with vegetables contributing the largest amount of dietary nitrates per serving [2, 9, 10]. Nitrates are inherently present in all plant materials, especially vegetables and forage crops, and accumulate when the plant matures in a nitrate rich environment [11]. Nitrates in drinking water are often the result of contamination of ground water by fertilizer and animal or human waste [2]. Cured meats, baked goods and cereals are notable sources of nitrite [12]. Nitrite salts are added to meats, poultry, and fish in minute quantities as a means of preservation; this has been a common practice for many centuries [13]. The endogenous conversion of nitrate to nitrite is also a significant source of exposure to nitrites; approximately 5% of ingested nitrates in food and water are converted to nitrite in the saliva [14]. Humans are exposed to *N*-nitroso compounds from exogenous sources and through endogenous formation. Dietary sources of nitrosamines include cured meats, beer, and smoked fish; these foods may contain preformed nitrosamines as the result of cooking and/or preservation methods [13, 15, 16]. Non-dietary sources include tobacco products, cosmetics, and occupational exposures in rubber or rocket fuel factories and leather tanneries [17-21].

As part of an ongoing study of the relation between prenatal exposure to these compounds and selected congenital malformations in offspring, we developed estimates of

nitrates, nitrites, and nitrosamines for food items in the Short Willet Food Frequency Questionnaire (WFFQ) [22, 23] as adapted for the National Birth Defects Prevention Study (NBDPS) [24]. Although an extensive body of literature exists on estimates of these compounds in foods, the extant data varies in quality, quantified estimates, country of origin and relevance to the U.S. population. Much of the literature was published before 1970 and may not accurately reflect current technological advances in laboratory methods and food preservation techniques. Furthermore, published estimates of nitrates, nitrites, and nitrosamines are not available for the WFFQ, a food frequency questionnaire that is commonly used in research studies.

## **Methods**

The NBDPS study collects dietary information using the adapted WFFQ [22, 23]. This adaptation was designed to make the questionnaire more suitable to the ethnic and racially diverse NBDPS study population and includes additional food items such as avocados, raw chili peppers, salsa, tortillas, cantaloupe, and refried beans.

Four steps were required to generate estimates of nitrates, nitrites, and nitrosamines for food items in the WFFQ: data collection, creation of a food database, selecting relevant values from that food database for our study population, and finally generating a summary estimate for each food item. To generate estimates, it was necessary to first develop a food database containing all relevant information associated with each published value (mean, range, cooking method, country, author, year, etc.) obtained from our literature search. This food database was consulted to identify appropriate values for our population based on our ranking system which accounted for the year(s) of sample collection and country of origin. Finally, for each food item we calculated a summary estimate based on the relevant values selected from the food database.

### **Step 1: Data Collection**

A thorough literature review was conducted to identify and categorize relevant reported values of nitrate, nitrite, and nitrosamine in foods. Multiple reference databases were searched for published literature reflecting nitrate, nitrite, and nitrosamine values in foods and alcoholic beverages. Reference databases queried included Medline, ISI Web of Science, Agricola, and Google Scholar. Each reference database was searched using the following terms: “nitrates,” “nitrites,” “nitrosamines,” “food content,” “dietary sources,” and combinations of the listed terms. Relevant published literature was reviewed but only sources reporting results for the food items listed on the WFFQ as adapted for the NBDPS were selected for inclusion [22-24]. For alcoholic beverages (beer, wine, liquor, and malt beverages) and for food items that returned no values for nitrates, nitrites, or nitrosamines an additional literature review was conducted. Food item estimates that were not available in the literature were substituted with similar food items with reported values. For example, a nitrosamine value for “orange juice” was not located using the search criteria previously mentioned; therefore, the value for “oranges” was used as a comparable substitution. There were also situations where some of the missing food items required recipes to generate an appropriate estimate. For example, a nitrate value for “salsa” was calculated using values for “onions”, “peppers”, and “tomatoes” in the proportions designated in a commonly available cookbook [25].

## **Step 2: Database Creation**

The food database was created as part of the process to generate summary estimates that are used to determine dietary intake of these compounds for U.S. women participants (giving birth during 1997 – 2005) in the NBDPS. The identified values were prioritized within each food item according to country of origin and the year of sample collection. The year of sample collection was used in order to select the time period that most accurately reflected the values of

these compounds in foods consumed by the NBDPS study population. However, the sample year was not always available; in those situations the year the article was received by the journal was used. When the above information was unavailable, we defaulted to the year of publication. Studies have shown that nitrate, nitrite, and nitrosamine content in food products likely varies over time and by region [26, 27]. To develop the most accurate and relevant values in terms of time and place for the NBDPS study population, priority was given to estimates from Western countries from 1980 to present. The ranks ranged from one (highest) to five (lowest) and are as follows: 1) U.S. and Canada 1980 to present; 2) countries with predominantly Western diets 1980 to present; 3) U.S. and Canada 1970-1979; 4) countries with predominantly Western diets 1970-1979; and 5) countries with predominantly non-Western diets. Thirty-four articles and four government reports were selected for inclusion in the food database and are listed in Table 1. As available, information was abstracted regarding cooking or preservation methods, methods of laboratory analysis, number of observations in each sample, ranges of values, means, and standard deviations. Although not specifically published in this article, the food database is available upon request.

### **Step 3: Value Selection**

Values compiled in the food database from the United States and Canada (rank 1) and other countries with traditionally Western diets (rank 2) from 1980 to present were automatically used to calculate the estimates for nitrates, nitrites, and nitrosamines in each food item. When there were fewer than five values reported for a food item ranked 1 or 2, values from the United States and Canada from 1970-1979 were used. Less relevant values with respect to country and year were used when no other estimates were available.

### **Step 4: Estimate Generation**

Table 1 lists the 30 references that met the value selection criteria for generating summary estimates of nitrates, nitrites, and nitrosamines in food items. A summary estimate for each food item was generated through a multi-step process using the mean or midpoint and the sample sizes from each study that met our selection criteria. A systematic approach to calculate these estimates was developed using STATA 10 [28]. A summary estimate for each food item was calculated as the weighted mean of the published values for that food item identified from the food database. The weights were calculated as the square root of the reported sample size for each value used to formulate the summary estimate. We would have preferred to use the inverse of the variance to weight the values but many of the reported values did not include a variance estimate. This process was implemented to create weighted total estimates that were representative of the reported values. The estimates are based on reported values selected to represent the U.S. food supply and potentially represent the exposure of the U.S. population to these compounds. This procedure was repeated for each food item.

The WFFQ as adapted for the NBDPS asks about dietary intake of 64 food items, some of which are categories of foods; however, we developed a single summary estimate per compound (nitrate, nitrite, and nitrosamines) for each food item. The process was straightforward in the case of individual food items, but required an additional procedure for category items. For example, the “processed meats” item contains component estimates for sausage, salami, lunchmeat, pâté, and preserved meat. To generate a single estimate for “processed meats”, weighted estimates for each individual component of that group were calculated. The summary estimate for “processed meats” was calculated by weighting the component estimates equally. The summary estimates were reported in mg/100g for nitrates and nitrites and µg/100g for nitrosamines.

For each food item, nitrates, nitrites, and nitrosamines were calculated for a typical serving using the summary estimates and the standard serving size in grams as defined by the NBDPS Nutrient Calculator when available [29]. For cereal, biscuits, fruit drinks, soy milk and tofu, the standard serving sizes were obtained using nutrition information provided on the label of commonly available products from a local grocery store.

The NBDPS Mother Questionnaire collects information regarding intake of alcoholic beverages by frequency and type of alcohol consumed. The questionnaire divides alcoholic beverages into five broad categories identified as beer, wine, mixed drinks, shot liquor, and other alcohol. We used standard serving sizes of alcohol in the United States as reported by the Centers for Disease Control: 12 ounces of beer, 8 ounces of malt liquor, 5 ounces of wine, or 1.5 ounces of liquor [30]. Based on the standard measures of alcohol, the serving size in grams for each of the categories was measured using commonly available samples of each type of beverage. Because estimates of nitrates and nitrites in alcoholic beverages were rarely reported in the literature, these compounds were not estimated for standard servings of these beverages. However, nitrosamine values were available and were used to calculate the estimated amount per serving using the same procedures used in the food calculations.

## **Results**

Table 2 displays the calculated amount per serving for food items grouped into dairy products; fruit; grains; meat and beans; vegetables; fats, oils, nuts, and sweets; and alcoholic beverages. The estimates in Table 2 are based on standard serving sizes for adults. Nitrate values ranged from 0 – 188.999 mg/serving with the highest concentrations occurring in vegetable products. Spinach and squash contain the highest amounts of nitrate per serving with

values of 188.999 and 43.608 mg respectively. Sweets, nuts, fats and oils contain very little nitrates per serving.

Nitrite values range from 0 - 1.840 mg/serving with the highest concentrations occurring in meat and bean products. Beef, pork, lamb, or *cabrito* (goat meat) as a main dish and liver contain the highest amounts of dietary nitrite per serving with 1.840 and 1.608 mg respectively. Negligible sources of dietary nitrite are found in cottage cheese, fats such as butter or margarine, and various fruit juices.

Nitrosamine values from food items ranged from 0 - 0.453  $\mu\text{g}$ /serving with the highest concentrations occurring in meat and dairy products. Beef, pork, lamb, or *cabrito* as a sandwich or main dish contains 0.324 and 0.453  $\mu\text{g}$ /serving. Cottage or ricotta cheese, fish and bacon contain high levels of nitrosamines with 0.266, 0.222 and 0.219  $\mu\text{g}$ /serving respectively. Alcoholic beverages also contain high levels of nitrosamines, with beer and malt beverages containing the highest amount of nitrosamines per serving at 0.531  $\mu\text{g}$  and 0.301  $\mu\text{g}$  respectively. Wine and liquor contain relatively little nitrosamines per serving with values of 0.019 and 0.027  $\mu\text{g}$  respectively. Fruits, vegetables, sweets, and fats do not contain significant amounts of nitrosamines per serving.

Table 3 displays the estimated amounts of nitrates, nitrites, and nitrosamines per 100 g of food item or alcoholic beverage. The food items are grouped into categories consisting of dairy products; fruit; grains; meat and beans; vegetables; fats, oils, nuts, and sweets; and alcoholic beverages. The values reported in Table 3 can be used to calculate amounts per serving for serving sizes other than those reported in Table 2.

## **Discussion**

Based on our estimates, total dietary nitrate intake per serving is most heavily influenced by vegetable consumption, specifically the green leafy varieties. However, nitrate intake in conjunction with vitamin C and possibly vitamin E may inhibit endogenous nitrosamine formation [31]. The benefits of fruit and vegetable consumption with their vitamin content most likely negate the potential harmful effects of nitrate intake from these sources. Our estimates indicate that nitrites and nitrosamines are most commonly associated with the consumption of meats, processed meats and fish. Based on our results, beer had the largest estimated amount of nitrosamine per serving.

Potential limitations in our study include sources of error associated with food item substitutions; different methodologies for measuring nitrates, nitrites, and nitrosamines in food; estimate calculations; and changes in concentration of these compounds in foods over time. The variability of nitrates, nitrites, and nitrosamines in food items may also be a source of error. Additionally, potential error stems from our substitutions and recipes used when the exact value for a food item was not available in the literature, although we expect the impact of these substitutions to be negligible. It is possible that nitrate, nitrite, and nitrosamine content in food items may vary by period measured, preparation, and analytical techniques used. To generate the most accurate estimate possible, we weighted individual values by their sample size. In most cases, weighting the values by the sample size more accurately reflects the available data. However, in a few cases, the estimate generated may be less accurate because of the over representation of values with extremely large samples sizes. Our study was also limited based on the relatively small number of references that were relevant to the current U.S. population based on year of publication and country of origin. Of the 30 references used to calculate estimates, only 11 were published since 1990 and none of those were based on U.S. or Canadian values.

Therefore, estimates of nitrates, nitrites, and nitrosamines should be updated with the appropriate analyses of food products currently available in the United States. Consideration should also be given to regional and ethnic differences in food availability and dietary intake.

Despite these limitations, our study also has several strengths. The estimates we generated are unique in that they were created to be used with the WFFQ, a commonly used research tool to assess dietary intake in research studies. Intake of nitrates, nitrites, and nitrosamines has likely changed over time because of procedural changes in food preservation and dietary behavioral changes [32]. Articles in the literature, such as those authored by Jakszyn *et al.* and Cornee *et al.*, provide values of nitrate, nitrite and nitrosamines in foods from the global literature; however, we were unable to locate summary estimates for the U.S. population that systematically accounted for time and country of origin [10, 33]. In the present study, we compiled, ranked, selected, and weighted the reported values to generate an estimate for each food item. Our estimates are not intended to be comprehensive in terms of all published literature regarding nitrate, nitrite, and nitrosamines in foods. Rather, the estimates were generated to reflect relevant information in the published literature that was representative of the U.S. population with respect to time and country of origin.

### **Conclusions**

Laboratory analysis of nitrate, nitrite, and nitrosamine content in food items commonly available in the United States should be conducted to determine current exposures to the U.S. population. In the absence of current data based on laboratory analysis, our estimates can be used to assess dietary exposures to these compounds. To the best of our knowledge, our estimates are the only ones published to date that are developed specifically for use with the WFFQ and intended for use with the U.S. population. The estimates provided may also be

useful in other research studies, specifically in exploring the relation between intake of *N*-nitroso compounds and their precursors with health conditions such as cancer and adverse pregnancy outcomes.

### **Declaration of Competing Interests**

The authors declare that they have no competing interests.

### **Acknowledgements**

This research was supported by Award Number R01ES015634 from the National Institute of Environmental Health Sciences. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Environmental Health Sciences or the National Institutes of Health. We thank Dr. Thomas McDonald for his review of the manuscript.

## Literature Cited

1. Brender J, Olive JM, Felkner M, Suarez L, Marckwardt W, Hendricks KA: **Dietary nitrites and nitrates, nitrosatable drugs, and neural tube defects.** *Epidemiology* 2004, **15**(3):330-336.
2. Ward MH, Dekok T, Levallois P, Brender J, Gulis G, Nolan BT, VanDerslice J: **Workgroup report: Drinking-water nitrate and health--recent findings and research needs.** *Environ Health Perspect* 2005, **113**(11):1607-1614.
3. Longnecker MP, Daniels JL: **Environmental contaminants as etiologic factors for diabetes.** *Environ Health Perspect* 2001, **109**(Suppl 6):871-876.
4. Bloomfield RA, Welsch CW, Garner GB, Muhrer ME: **Effect of dietary nitrate on thyroid function.** *Science* 1961, **134**:1690.
5. Pomeranz A, Korzets Z, Vanunu D, Krystal H: **Elevated salt and nitrate levels in drinking water cause an increase in blood pressure in school children.** *Kidney Blood Press Res* 2000, **23**(6):400-403.
6. Ivankovic S: **Teratogenic and carcinogenic effects of some chemicals during prenatal life in rats.** *Natl Canc Inst Monogr* 1979(51):103-115.
7. Inouye M, Murakami U: **Teratogenic effect of N-methyl-N-nitro-N-nitrosoguanidine in mice.** *Teratology* 1978, **18**(2):263-267.
8. Koyama T, Handa H, Matsumoto S: **Methylnitrosourea-induced malformations of the brain in SD-JCL rat.** *Arch Neurol* 1970, **22**:342-347.
9. Walters CL: **Nitrate and nitrite in foods.** New York: Ellis Horwood; 1991.
10. Jakszyn P, Agudo A, Ibanez R, Garcia-Closas R, Pera G, Amiano P, Gonzalez CA: **Development of a food database of nitrosamines, heterocyclic amines, and polycyclic aromatic hydrocarbons.** *J Nutr* 2004, **134**(8):2011-2014.
11. Ashton MR: **The occurrence of nitrates and nitrites in food.** *The British Food Manufacturing Industries Research Association: Literature Survey No 7* 1970:1-32.
12. National Academy of Sciences: **Nitrate, nitrite, and nitrogen oxides: environmental distribution and exposure of humans.** In. Washington D.C.: National Academies Press; 1981a: 3-78.
13. Hotchkiss JH: **A review of the current literature on N-nitroso compounds in foods.** *Adv Food Res* 1987, **31**:53-115.

14. Choi BC: **N -nitroso compounds and human cancer: a molecular epidemiologic approach.** *Am J Epidemiol* 1985, **121**(5):737-743.
15. Lijinsky W: **N -nitroso compounds in the diet.** *Mutat Res* 1999, **443**(1-2):129-138.
16. Tricker AR, Preussmann R: **Volatile and nonvolatile nitrosamines in beer.** *J Cancer Res Clin Oncol* 1991, **117**:130-132.
17. Stepanov I, Hecht SS: **Tobacco-specific nitrosamines and their pyridine- N -glucuronides in the urine of smokers and smokeless tobacco users.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:885-891.
18. Bernert JT, Jain RB, Prikle JL, Wang L, Miller BB, Sampson EJ: **Urinary tobacco-specific nitrosamines and 4-aminobiphenyl hemoglobin adducts measured in smokers of either regular or light cigarettes.** *Nicotine Tob Res* 2005, **7**(5):729-738.
19. Havery DC, Chou HJ: **N-nitrosamines in cosmetic products.** *Cosmet Toiletries* 1994, **109**:53-61.
20. Walters KA, Brain KR, Dressler WE, Green DM, Howes D, James VJ, Kelling CK, Watkinson AC, Gettings SD: **Percutaneous penetration of N-nitroso-N-methyldodecylamine through human skin in vitro: application from cosmetic vehicles.** *Food Chem Toxicol* 1997, **35**(7):705-712.
21. National Academy of Sciences: **N-nitroso compounds: environmental distribution and exposure in humans.** In: *In: The Health Effects of Nitrate, Nitrite, and N-nitroso compounds.* Washington D.C.: National Academies Press; 1981b: 3-51.
22. Willett WC, Reynolds RD, Cottrell-Hoehner S, Sampson L, Brown ML: **Validation of a semi-quantitative food frequency questionnaire: comparison with a 1-year diet record.** *J Am Diet Assoc* 1987, **87**:43-47.
23. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE: **Reproducibility and validity of a semiquantitative food frequency questionnaire.** *Am J Epidemiol* 1985, **122**:51-65.
24. Yoon PW, Rasmussen SA, Lynberg MC, Moore CA, Anderka M, Carmicheael SL, Costa P, Druschel C, Hobbs CA, Romitti PA *et al*: **The National Birth Defects Prevention Study.** *Public Health Rep* 2001, **166**(Suppl 1):32-40.
25. Westmoreland S, The Good Housekeeping Institute: **Good Housekeeping Step by Step Cookbook.** New York: Hearst Books; 2003.
26. Paik DC, Saborio DV, Oropeza R, Freeman HP: **The epidemiological enigma of gastric cancer rates in the US: was grandmother's sausage the cause?** *Int J Epidemiol* 2001, **30**(1):181-182.

27. Binkerd EF, Kolari OE: **The history and use of nitrate and nitrite in the curing of meat.** *Food Cosmet Toxicol* 1975, **13**(6):655-661.
28. StataCorp: **Stata Statistical Software: Release 10.** In. College Station, TX: StataCorp LP; 2007.
29. Farrell J: **Draft NBDPS Nutrient Calculations, Version 3 of the Analytic Database.** In.; 2004.
30. **Quick Stats: General information on alcohol use and health**  
[[http://www.cdc.gov/alcohol/quickstats/general\\_info.htm](http://www.cdc.gov/alcohol/quickstats/general_info.htm)]
31. Bartsch H, Calmels S, Ohshima H, Pignatelli B: **Inhibition of nitrosation.** *Basic Life Sci* 1993, **61**:27-44.
32. Dietrich M, Block G, Pogoda JM, Buffler P, Hecht SS, Preston-Martin S: **A review: dietary and endogenously formed N-nitroso compounds and risk of childhood brain tumors.** *Cancer Causes Control* 2005, **16**(6):619-635.
33. Cornee J, Lairon D, Velema J, Guyader M, Berthezene P: **An estimate of nitrate, nitrite, and N-nitrosodimethylamine concentrations in French food products or food groups.** *Sciences des Aliments* 1992, **12**(2):155-197.
34. Canas BJ, Havery DC, Joe FL, Fazio T: **Current trends in levels of volatile N-nitrosamines in fried bacon and fried-out bacon fat.** *J Assoc of Anal Chem* 1986, **69**(6):1020-1021.
35. Chung SY, Kim JS, Kim M, Hong MK, Lee JO, Kim CM, Song IS: **Survey of nitrate and nitrite contents of vegetables grown in Korea.** *Food Addit Contam* 2003, **20**(7):621-628.
36. Dennis MJ, Cripps GS, Tricker AR, Massey RC, McWeeny DJ: **N-nitroso compounds and polycyclic aromatic hydrocarbons in Icelandic smoked cured mutton.** *Food Chem Toxicol* 1984, **22**:305-306.
37. Food Standards Agency: Joint Food Safety Standards Group: **Food Surveillance Information Sheet.** In.: Ministry of Agriculture, Fisheries, and Food United Kingdom; 1998.
38. French National Inventory: **Ministeres de l'Environnement, de l'Agriculture, de la Recherche et de la Technologie, de l'Industrue et de la Sante, Paris.** In.; 1982.
39. Fudge R, Truman RW: **The nitrate and nitrite contents of meat products: a survey by public analysts' laboratories in South Wales and the South West of England.** *J Assoc Publ Anal* 1973, **11**:19-27.
40. Havery DC, Hotchkiss JH, Fazio T: **Rapid determination of volatile N-nitrosamines in nonfat dry milk.** *J Dairy Sci* 1982, **65**:182-185.

41. Howe GR, Harrison L, Jain M: **A short diet history for assessing dietary exposure to N-nitrosamines in epidemiologic studies.** *Am J Epidemiol* 1986, **124**:595-602.
42. Huang DP, Ho JH, Webb KS, Wood BJ, Gough TA: **Volatile nitrosamines in salt-preserved fish before and after cooking.** *Food Cosmet Toxicol* 1981, **19**:167-171.
43. Key PE, Baylor JP, Massey RC, McWeeny DJ: **Nitrosodimethylamine levels in fish cooked by natural gas and by electricity.** *Food Technol* 1982, **17**:703-708.
44. Klein D, Girard AM, Cabarro C, Debry G: **Presence of volatile nitrosamines in food.** *Ann Nutr Aliment* 1980, **34**:915-928.
45. Knight TM, Forman D, Al-Dabbagh SA, Doll R: **Estimation of dietary intake of nitrate and nitrite in Great Britain.** *Food Chem Toxicol* 1987, **25**(4):277-285.
46. Kyriakidis NB, Tarantili-Georgiou K, Tsani-Batzaka E: **Nitrate and Nitrite Content of Greek Cheeses.** *J Food Comp Anal* 1997, **10**(4):343-349.
47. Lakritz L, Pensabene JW: **Survey of fluid and nonfat dry milks for N-nitrosamines.** *J Dairy Sci* 1981, **64**:371-374.
48. Libbey LM, Scanlan RA, Barbour JF: **N-Nitrosodimethylamine in dried dairy products.** *Food Cosmet Toxicol* 1980, **18**:459-461.
49. Mahieu H, Luquet FM, Mouillet L, Boudier JF: **Occurrence of nitrates and nitrites in milk, Evaluation of the nitrate and nitrite content of French milk; 1973, 1974, 1975, 1979 surveys.** *Ann Nutr Aliment* 1980, **34**:1045-1052.
50. Maki T, Tamura Y, Shimamura Y, Naoi Y: **Estimate of the volatile nitrosamine content of Japanese food.** *Bull Environ Contam Toxicol* 1980, **25**:257-261.
51. Matsui M, Ohshima H, Kawabata T: **Increase in the nitrosamine content of several fish products upon broiling.** *Bull Jap Soc Sci Fish* 1980, **46**:587-590.
52. Merino L, Darnerud PO, Edberg U, Aman P, Castillo MDP: **Levels of nitrate in Swedish lettuce and spinach over the past 10 years.** *Food Addit Contam* 2006, **23**(12):1283-1289.
53. Osterdahl BG: **Volatile nitrosamines in foods on the Swedish market and estimation of their daily intake.** *Food Addit Contam* 1988, **5**:587-595.
54. Pedersen E, Meyland I: **Nitrate, nitrite, and volatile nitrosamines in pickled fish prepared with addition of nitrate.** *Z Lebensm Unters Forsch* 1981, **173**:359-361.
55. Petersen A, Stoltze S: **Nitrate and nitrite in vegetables on the Danish market: content and intake.** *Food Addit Contam* 1999, **16**(7):291-299.

56. Pobel D, Riboli E, Cornee J, Hemon B, Guyader M: **Nitrosamine, nitrate and nitrite in relation to gastric cancer: A case-control study in Marseille, France.** *Eur J Epidemiol* 1995, **11**:67-73.
57. Saccani G, Tanzi E: **Determination of nitrite, nitrate, and glucose-6-phosphate in muscle tissues and cured meat by IC/MS.** *J AOAC Int* 2006, **89**(3):712-719.
58. Scanlan RA, Barbour JF, Hotchkiss JH, Libbey LM: **N-nitrosodimethyl-amine in beer.** *Food Cosmet Toxicol* 1980, **18**:27-29.
59. Sen NP, Seaman S, McPherson M: **Nitrosamines in alcoholic beverages.** *J Food Saf* 1980, **2**:13-18.
60. Sen NP, Seaman S, Baddoo PA, Weber D: **Further studies on the formation of nitrosamines in cured pork products packaged in elastic rubber nettings.** *J Food Sci* 1988, **53**:731-738.
61. Siciliano J, Krulick S, Heisler EG, Schwartz JH, White JW: **Nitrate and Nitrite Content of Some Fresh and Processed Market Vegetables.** *J Agric Food Chem* 1975, **23**(3):461-464.
62. Siddiqi M, Tricker AR, Preussmann R: **The occurrence of preformed N-nitroso compounds in food samples from a high risk area of esophageal cancer in Kashmir, India.** *Cancer Lett* 1988, **39**:37-43.
63. Spiegelhalter B, Eisenbrad G, Preussmann R: **Volatile nitrosamines in food.** *Oncology* 1980, **37**:211-216.
64. Tamme T, Reinik M, Roasto M, Juhkam K, Tenno T, Kiis A: **Nitrates and nitrites in vegetables and vegetable-based products and their intakes by the Estonian population.** *Food Addit Contam* 2006, **23**(4):355-361.
65. Thomson B, Swallow B: **Nitrates and nitrites dietary exposure and risk assessment.** In. Christchurch, New Zealand; 2004: 1-37.
66. Tricker AR, Pfundstein B, Theobald E, Preussmann R, Spiegelhalter B: **Mean daily intake of volatile N-nitrosamines from foods and beverages in West Germany in 1989-1990.** *Food Chem Toxicol* 1991, **29**:729-732.
67. Vecchio AJ, Hotchkiss JH, Bisogni CA: **N-nitrosamine Ingestion from consumer cooked bacon.** *J Food Sci* 1986, **51**:754-756.
68. VonCollett P: **Beitrag zur bestimmung von Nitrit und Nitrat in lebensmitteln.** *Deut Lebensm-Rundsch* 1983, **79**(11):370-375.
69. Yamamoto M, Iwata R, Ishiwata H, Yamada T, Tanimura A: **Determination of volatile nitrosamine levels in foods and estimation of their daily intake in Japan.** *Food Chem Toxicol* 1984, **22**:61-64.

**Table 1.** References used in food database and estimate generation

Citation	Year(s) Values Collected <sup>a</sup>	Country <sup>b</sup>	Rank <sup>c</sup>	Included in Estimate <sup>d</sup>
Canas, et al., 1986 [34]	1985	US	1	√
Chung, et al., 2003 [35]	1998	KR	5	√
Cornee, et al., 1992 [33]	1992	FR	2	√
Dennis, et al., 1984 [36]	1983	IS	5	-
Food Standards Agency, 1998 [37]	1997	UK	2	√
French National Inventory, 1982 [38]	1982	FR	2	√
Fudge and Truman, 1973 [39]	1973	UK	4	-
Havery, et al., 1982 [40]	1981	US	1	√
Howe, et al., 1986 [41]	1985	CA	1	√
Huang, et al., 1981 [42]	1980	HK	5	-
Key, et al., 1982 [43]	1982	UK	2	√
Klein, et al., 1980 [44]	1980	FR	2	√
Knight, et al., 1987 [45]	1973-1978	UK	4	-
Kyriakidis, et al., 1997 [46]	1995	GR	5	-
Lakritz and Pensabene, 1981 [47]	1980	US	1	√
Libbey, et al., 1980 [48]	1979	US	3	√
Mahieu, et al., 1980 [49]	1980	FR	2	√
Maki, et al., 1980 [50]	1976-1980	JP	5	√
Matsui, et al., 1980 [51]	1979	JP	5	-
Merino, et al., 2006 [52]	1995-2005	SE	2	√
National Academy of Sciences, 1981a [12]	1975-1979	US	3	√

Osterdahl, 1988 [53]	1980-1986	SE	2	√
Pedersen and Meyland, 1981 [54]	1981	DE	2	√
Petersen and Stoltze, 1999 [55]	1993-1997	DK	2	√
Pobel, et al., 1995 [56]	1994	FR	2	√
Saccani and Tanzi, 2006 [57]	2005	IT	2	√
Scanlan, et al., 1980 [58]	1979	US	3	-
Sen, et al., 1980 [59]	1979	CA	3	√
Sen, et al., 1988 [60]	1988	CA	1	√
Siciliano, et al., 1975 [61]	1973-1974	US	3	√
Siddiqi, et al., 1988 [62]	1988	IN	5	-
Spiegelhalder, et al., 1980 [63]	1980	DE	2	√
Tamme, et al., 2006 [64]	2003-2004	EE	5	√
Thomson and Swallow, 2004 [65]	2003	NZ	2	√
Tricker, et al., 1991 [66]	1989-1990	DE	2	√
Vecchio, et al., 1986 [67]	1984	US	1	√
VonCollett, 1983 [68]	1983	DE	2	√
Yamamoto, et al., 1984 [69]	1982	JP	5	√

<sup>a</sup>Year(s) sample collected is reported, when not available year received by journal is used and year of publication is used as when all other dates are unavailable

<sup>b</sup>CA = Canada, DE = Germany, DK = Denmark, EE = Estonia, FR = France, GR = Greece, HK = Hong Kong, IN = India, IS = Iceland, IT = Italy, JP = Japan, KR = South Korea, NZ = New Zealand, SE = Sweden, UK = United Kingdom, US = United States

<sup>c</sup>Rank 1= U.S. and Canada 1980 to present; Rank 2= countries with predominantly Western diets 1980 to present; Rank 3= U.S. and Canada 1970-1979; Rank 4= countries with predominantly Western diets 1970-1979; and Rank 5= countries with predominantly non-Western diets

<sup>d</sup>(√) = included in estimate; (-) = excluded from estimate

**Table 2.** Estimates of nitrates, nitrites and nitrosamines in food items and alcoholic beverages by the National Birth Defects Prevention Study calculated serving size

<b>Food Item</b>	<b>Serving Size g</b>	<b>Nitrates mg/serving</b>	<b>Nitrites mg/serving</b>	<b>Nitrosamines µg/serving</b>
<i><b>Dairy Products</b></i>				
Cheese (1 slice or 1 oz)	28	0.400	0.023	0.066
Cottage or Ricotta cheese (1/2 cup)	113	0.260	0.000	0.266
Ice cream (1/2 cup)	72	0.209	0.038	0.031
Skim or low fat milk (8 oz glass)	244	0.862	0.011	0.209
Whole Milk (8 oz glass)	244	0.212	0.001	0.065
Yogurt ( 1 cup)	245	0.333	0.071	0.002
<i><b>Fruit Products</b></i>				
Avocado (1) or guacamole (1 cup)	201	5.278	0.042	0.010
Bananas (1)	114	2.280	0.043	0.006
Cantaloupe (1/4 melon)	134	12.730	0.051	0.006
Fresh apples or pears (1)	138	1.436	0.014	0.007
Hawaiian Punch, lemonade, or other fruit drinks (1 glass)	244	2.367	0.000	0.000
Orange juice (small glass)	186	3.720	0.000	0.000
Oranges (1)	184	3.680	0.000	0.000
Other fruits fresh, frozen, or canned (1/2 cup)	142	3.728	0.030	0.007
Peaches, apricots, plums, or nectarines (1	87	0.564	0.014	0.004

or 1/2 cup)

Raw chile peppers, jalapeño (1)	14	0.742	0.009	0.000
Salsa (1 cup) (fruit or tomato)	259	9.094	0.140	0.000
Tomatoes (1) or tomato juice (small glass)	123	3.953	0.054	0.000

***Grain Products***

Biscuit, scone, croissant and muffin (1)	80	0.328	0.032	0.012
Cereal	30	0.138	0.039	0.012
Dark bread (slice) including wheat pita bread	25	0.213	0.030	0.000
Rice or pasta (1 cup)	140	2.305	0.303	0.000
Tortilla (1)	19	0.304	0.031	0.002
White bread (slice), pita bread, bagels and crackers	25	0.410	0.042	0.002

***Meat and Bean Products***

Bacon (2 slices)	16	1.449	0.467	0.219
Beans or lentils, baked or dried (1/2 cup)	131	1.179	0.341	0.000
Beans, Refried (1 cup)	252	2.268	0.655	0.000
Beef, pork, lamb or cabrito as a main dish (4-6 oz)	140	8.188	1.840	0.453
Beef, pork, lamb or cabrito sandwich or mixed dish	100	5.849	1.314	0.324
Chicken Livers (1 oz)	28	0.958	0.571	0.007
Chicken or Turkey (4-6 oz)	88	0.583	0.478	0.086

Eggs (1)	58	0.311	0.095	0.000
Fish (3-6 oz)	112	1.027	0.349	0.222
Hamburger (1 patty)	77	6.102	0.193	0.071
Hot dogs (1)	45	11.565	1.220	0.128
Liver (3-4 oz)	82	10.325	1.608	0.022
Organ meats and tongue (3-4 oz)	85	0.882	0.480	0.062
Peas or lima beans (1/2 cup frozen, canned)	80	2.634	0.070	0.000
Processed meats - sausage, salami, lunchmeat, pâté, and preserved meats (piece or slice)	28	1.700	0.314	0.124
<b><i>Vegetables Products</i></b>				
Broccoli (1/2 cup)	78	26.607	0.251	0.000
Cabbage, cauliflower, brussel sprouts (1/2 cup)	75	12.976	0.143	0.000
Carrots, cooked (1/2 cup)	78	13.013	0.115	0.000
Carrots, raw (1/2 cup or 2-4 sticks)	36	6.006	0.053	0.000
Corn (1 ear or 1/2 cup frozen, canned)	82	3.690	0.164	0.000
Potatoes baked, boiled (1) or mashed (1 cup)	156	22.673	0.164	0.000
Soy milk or soy yogurt (8 oz)	244	13.333	0.220	0.000
Spinach or collard greens, cooked (1/2 cup)	90	188.999	0.177	0.000

Squash (1/2 cup)	103	43.608	0.082	0.000
String Beans (1/2 cup)	68	11.767	0.128	0.000
Tofu, tempeh or soy burgers (4oz)	113	6.174	0.102	0.000
Yams or sweet potatoes (1/2 cup)	68	3.128	0.048	0.000

***Fats, Oils, Nuts, and Sweets***

Butter (pat), added to food or bread	5	0.000	0.000	0.000
Cake (slice) or donut (1)	64	1.600	0.083	0.000
Candy without chocolate (1 oz)	28	0.000	0.000	0.010
Chocolate (1 oz)	21	0.000	0.000	0.010
Cookies (1)	25	0.092	0.011	0.002
French fried potatoes (4 oz)	112	4.144	0.078	0.000
Margarine (pat), added to food or bread	5	0.002	0.000	0.000
Nuts (small packet or 1 oz)	18	0.104	0.006	0.000
Oil and Vinegar dressing (1 tbs)	15	0.000	0.000	0.000
Peanut butter (1 tbs)	16	0.093	0.006	0.000
Pie (slice)	125	2.338	0.138	0.000
Potato or corn chips (small bag or 1oz)	28	1.049	0.020	0.000

***Alcoholic Beverages***

Beer (12 fl. oz/354.9 ml)	357	-	-	0.531
Wine (5fl. oz/147.9 ml)	136	-	-	0.019
Liquor and mixed drinks (1.5 fl. oz/44.4 ml)	41	-	-	0.027
Malt beverages (8 fl. oz/236.6 ml)	245	-	-	0.301

**Table 3.** Estimates of nitrates, nitrites and nitrosamines per 100 grams of food item or alcoholic beverage

<b>Food Item</b>	<b>Nitrates mg/100g</b>	<b>Nitrites mg/100g</b>	<b>Nitrosamines µg/100g</b>
<i><b>Dairy Products</b></i>			
Cheese	1.428	0.083	0.235
Cottage or Ricotta cheese	0.230	0.000	0.235
Ice cream	0.291	0.053	0.043
Skim or low fat milk	0.353	0.004	0.086
Whole Milk	0.087	0.000	0.027
Yogurt	0.136	0.029	0.001
<i><b>Fruit and Related Products</b></i>			
Avocado or guacamole	2.626	0.021	0.005
Bananas	2.000	0.038	0.005
Cantaloupe	9.500	0.038	0.005
Fresh apples or pears	1.040	0.010	0.005
Hawaiian Punch, lemonade, or other fruit drinks	0.970	0.000	0.000
Orange juice	2.000	0.000	0.000
Oranges	2.000	0.000	0.000
Other fruits fresh, frozen, or canned	2.626	0.021	0.005
Peaches, apricots, plums, or nectarines	0.648	0.016	0.005
Raw chile peppers, jalapeño	5.300	0.063	0.000
Salsa - fruit or tomato	3.510	0.054	0.000
Tomatoes or tomato juice	3.214	0.044	0.000

***Grain Products***

Biscuit, scone, croissant and muffin	0.410	0.040	0.015
Cereal	0.460	0.132	0.033
Dark bread - including wheat pita bread	0.850	0.120	0.000
Rice or pasta	1.647	0.217	0.000
Tortilla	1.640	0.166	0.009
White bread, pita bread, bagels and crackers	1.640	0.166	0.009

***Meat and Bean Products***

Bacon	9.058	2.918	1.369
Beans or lentils, baked or dried	0.900	0.260	0.000
Beans, Refried	0.900	0.260	0.000
Beef, pork, lamb or cabrito as a main dish	5.849	1.314	0.324
Beef, pork, lamb or cabrito as a sandwich or mixed dish	5.849	1.314	0.324
Chicken Livers	3.420	2.040	0.027
Chicken or Turkey	0.666	0.546	0.099
Eggs	0.536	0.164	0.000
Fish	0.917	0.312	0.198
Hamburger	7.900	0.250	0.092
Hot dogs	25.700	2.710	0.285
Liver	12.668	1.973	0.027
Organ meats and tongue	1.038	0.565	0.073
Peas or lima beans	3.293	0.088	0.000

Processed meats - sausage, salami, lunchmeat, pâté, and preserved meats	5.997	1.106	0.439
--	-------	-------	-------

***Vegetables and Related Products***

Broccoli	34.111	0.322	0.000
Cabbage, cauliflower, or brussel sprouts	17.301	0.190	0.000
Carrots, cooked	16.683	0.147	0.000
Carrots, raw	16.683	0.147	0.000
Corn	4.500	0.200	0.000
Potatoes baked, boiled or mashed	14.534	0.105	0.000
Soy milk or soy yogurt	5.464	0.090	0.000
Spinach or collard greens, cooked	209.999	0.196	0.000
Squash	42.544	0.080	0.000
String Beans	17.304	0.189	0.000
Tofu, tempeh or soy burgers	5.464	0.090	0.000
Yams or sweet potatoes	4.600	0.070	0.000

***Fats, Oils, Nuts, and Sweets***

Butter - added to food or bread	0.000	0.000	0.000
Cake or donut	2.500	0.130	0.000
Candy without chocolate	0.000	0.000	0.034
Chocolate	0.000	0.000	0.046
Cookies	0.370	0.043	0.007
French fried potatoes	3.700	0.070	0.000
Margarine - added to food or bread	0.050	0.000	0.000

Nuts	0.580	0.035	0.000
Oil and Vinegar dressing	0.000	0.000	0.000
Peanut butter	0.580	0.035	0.000
Pie	1.870	0.110	0.000
Potato or corn chips	3.700	0.070	0.000
<b><i>Alcoholic Beverages</i></b>			
Beer	-	-	0.149
Wine	-	-	0.014
Liquor and mixed drinks	-	-	0.059
Malt beverages	-	-	0.123