

# From functional food to medicinal product: Systematic approach in analysis of polyphenolics from propolis and wine

Marica Medić-Šarić \*<sup>1</sup>, Vesna Rastija<sup>2</sup>, Mirza Bojić<sup>1</sup> and Željko Maleš<sup>3</sup>

Address:

<sup>1</sup>Department of Medicinal Chemistry, Faculty of Pharmacy and Biochemistry,  
University of Zagreb, A. Kovačića 1, 10000 Zagreb, Croatia

<sup>2</sup>Faculty of Agriculture, Josip Juraj Strossmayer University of Osijek, Trg Sv.  
Trojstva 3, 31000 Osijek, Croatia

<sup>3</sup>Department of Pharmaceutical Botany, Faculty of Pharmacy and Biochemistry,  
University of Zagreb, Schrottova 39, 10000 Zagreb, Croatia

\*Corresponding author

Email:

Marica Medić-Šarić: [bebamms@pharma.hr](mailto:bebamms@pharma.hr)

Vesna Rastija: [vesna.rastija@pfos.hr](mailto:vesna.rastija@pfos.hr)

Mirza Bojić: [mbojic@pharma.hr](mailto:mbojic@pharma.hr)

Željko Maleš: [zmales@pharma.hr](mailto:zmales@pharma.hr)

## **Abstract**

In the last decade we have been working on standardization of propolis extract and determination of active constituents of wine those are rich in polyphenolics and have nutritional as well as therapeutic value. Here we are summarizing our results and providing overview on systematic approach how to analyse natural products rich in flavonoids and phenolic acids.

Chromatographic methods (thin layer chromatography and high performance liquid chromatography) were used for identification, quantification, and characterization of individual flavonoid or phenolic acid. Total content of active constituents and antioxidant activity were determined by spectrophotometry. Pharmacokinetic parameters were determined by high performance liquid chromatography and using appropriate software. Quantitative structure-activity relationship study of antioxidant activity was conducted, as well as assessment of prolonged propolis supplementation on antioxidative status of organism.

Thin layer chromatography-densitometry has been proven as quick and reliable method for standard analysis of propolis and wine; the best mobile phase being chloroform - methanol - formic acid (98-100%) in ratio 44 : 3.5 : 2.5 (v/v). Higher number of polyphenolics was determined by high performance liquid chromatography; 15 compared to 9 by thin layer chromatography. Interactions *in situ* with acetylsalicylic acid were detected with most of polyphenolics analysed. Plasma protein binding and blood-barrier penetration was greatest for flavone. The interactions with human serum albumin have been greater than 95% for all flavonoids analysed. The prolonged propolis consumption increased superoxide dismutase activity.

The necessity of standardization of natural products and their registration as functional nutraceuticals demand easy, quick and inexpensive methods of analysis. In this work we provided overview of analytical part for polyphenolics that could be used as data for possible registration of final products either as functional food or medicinal product.

This feature introduces the readers to the authors' research through a concise overview of the selected topic. Reference to important work from others in the field is included.

## **Background**

The main causes of death in western countries are cardiovascular diseases and cancer. Over 50 percent of the population has some kind of chronic condition (high blood pressure, high cholesterol, arthritis, diabetes, asthma, osteoporosis...), so the goal of the many researches in the last decades is improving the quality of life. Much of the interested was transferred to homeopathy, alternative, and folk medicine, evincing polyphenols as one of the main nutraceuticals [1].

The most abundant sources of polyphenols, mainly flavonoids and phenolic acids, are propolis and wine. Flavonoids and phenolic acids have antibacterial, antifungal, antiviral, antineoplastic, hepatoprotective, immunomodulating, and antiinflammatory properties. Their use has been proven beneficial in allergies, asthma, diabetes, hypertension, micro bleeding, etc. Much of these pharmacological effects can be associated with antioxidant activity. Hence, antioxidant activity is the most studied property of polyphenols [2].

In the last decade, we were working on several projects with the aim of developing standardized extracts of propolis and determination of active constituents of wine, transferring knowledge from the laboratory to everyday-practice for the purpose of

analyzing natural products. To accomplish this we used rather simple, reliable, and quick spectrophotometric and liquid chromatography methods for analysis of polyphenols, the results of which are reviewed here. These were used to suggest mixtures of propolis that will provide the best final product.

In this review we wanted to systematize the approach in analysis of polyphenols rich samples providing mostly unpublished data<sup>1</sup>. Some basic concepts of optimization of chromatographic system, chromatographic parameters describing pharmacokinetic and quantitative structure-activity relationship are described. The use of liquid chromatography for different objectives is thoroughly illustrated: identification, quantification, characterization; ADME study (Absorption, Distribution, Metabolism, Excretion) and interactions. To demonstrate *in vivo* antioxidant effects of polyphenols from propolis preliminary clinical trial of prolonged propolis is presented.

### **Flavonoids and Phenolic Acids. Propolis and Wine**

Flavonoids and phenolic acids are plant polyphenols characterized by having one or more phenolic hydroxyl groups or carboxylic group in phenolic acids. The major classes of polyphenols are phenolic acids (hydroxycinnamic and hydroxybenzoic acids), flavonoids, anthocyanins, and tannins (Figure 1) [2,3].

Propolis is a resinous substance collected by honeybees (*Apis mellifera*, L.) from a variety of plant sources. Its chemical composition is very complex and varies with geographic origin, depending on the local flora and phenology of the source plants. The benefits of effects of propolis on human health were recognized thousand of years ago. It has wide range of biological activities including antibacterial, antiviral,

---

<sup>1</sup> Published data is marked as reference.

antiinflammatory, and antioxidative [4]. To illustrate the regions where our propolis samples originate, the map of Croatia is provided on Figure 2.

Wine is an alcoholic beverage that contains a large amount of different polyphenols extracted from grapes during the processes of vinification. These molecules are responsible for color, acerbity, flavor, and antioxidant properties of wine [5].

### **Optimization of Chromatographic Systems [6]**

The purpose of optimization of chromatographic systems is to find the one that shows the greatest difference in identification characteristics between substances e.g. retention factor ( $R_F$ ) value in thin layer chromatography.

To accomplish this numerical taxonomy methods were used. Taxonomic entities are polyphenols and their numerical characteristic is  $R_F$  value. Matrix of resemblance of  $R_F$  values is analyzed to find the mobile phase in which the resemblance is the lowest and separation of analyzed substances is optimal. In short, the matrix of  $N$  substances and  $t$  mobile phases is reduced using different algorithms to form a cluster of similar mobile phases, the procedure is repeated until the final cluster including all mobile phases is formed and the results are presented in form of dendrogram.

Complementary mathematical methods are used. If the  $R_F$  values are distributed into groups of width  $E$ , in  $R_F$  units, then in group  $k$   $n_k R_F$  values are present, and the average information content is described by Shannon equation:

$$I(X) = - \sum_k \frac{n_k}{n} \log_2 \frac{n_k}{n}, \quad (1)$$

with the assumption that substances with the  $R_F$  value in the same group can not be identified. The higher the average information content is, the better results of separation for mobile phase are. To positively identify two substances difference between  $R_F$  values has to be greater than  $E$  (usually 0.05). Discriminating power ( $DP$ )

is used as a measure of the effectiveness of chromatographic systems. The  $DP$  of a chromatographic system is the probability of separating two flavonoids (or phenolic acids) selected at random from a specific substance population. Two flavonoids (or phenolic acids) are chromatographically similar if the differences in their identification values do not exceed the error factor  $E$ . In case of mixture of eluents  $DP$  is defined as probability that two randomly selected substances can be distinguished by one mobile phase. Mathematically it can be expressed by the following equation:

$$DP_k = 1 - \frac{2M}{N(N-1)}, \quad (2)$$

where  $k$  is the number of mobile phases,  $N$  number of substances analyzed and  $M$  the total number of matching pairs. The average number of similar substances  $T$ , for each mobile phase, is defined by expression:

$$T = 1 + \frac{N-1}{1- DP_k}. \quad (3)$$

The highest  $DP$  and the smallest  $T$  value are attributed to the most appropriate combination of eluents.

This procedure of optimization has been used in the analysis of numerous plant samples: *Zizyphus jujuba* Mill., *Chamomilla recutita* (L.) Rauschert, *Rosmarini folium*, *Guiera senegalensis* J.F.GMEL., *Rhamni cathartici fructus*, *Lavandulae flos*, *Sambuci flos*, *Helleborus atrorubens* Waldst. et Kit. etc. [7-14].

### **Chromatographic Parameters Describing Pharmacokinetics of Polyphenols [15]**

As our samples are consumed orally absorption and distribution of polyphenols are of interest. Absorption through the gastrointestinal tract (GIT) membrane, plasma protein binding and the penetration through blood brain barrier (BBB) was studied using *in silico* approaches and liquid chromatography.

In reverse phase thin layer chromatography (RP-TLC), the chromatographic parameter  $R_M$  is used to describe the lipophilicity of a substance:

$$R_M = \log \frac{b-a}{a} = \log \frac{1-R_F}{R_F} \quad (4)$$

It represents a logarithm of ratio of spot distance from the front and from the start. The higher the value of  $R_F$ , chromatographic parameter is lower and the substance is more polar. As  $R_M$  depends on the percentage of the organic modifier in mobile phase,  $R_{Mw}$  value with 0% of organic modifier is used and can be extrapolated using the equation:

$$R_M = R_{Mw} + b\phi \quad (5)$$

where  $b$  is the coefficient of the slope and  $\phi$  is the concentration of the organic modifier.

In reverse phase high performance liquid chromatography (RP-HPLC)  $\log k$  is used as a measurement of distribution between hydrophobic (stationary) and hydrophilic (mobile) phase:

$$\log k = \log \frac{C_{stationary}}{C_{mobile}} = \log \frac{t_R - t_0}{t_0} \quad (6)$$

The values of  $\log k$  are compared at a 0% concentration of organic modifier based on the equation:

$$\log k = \log k_w + b\phi \quad (7)$$

As mucosa membranes are modified with phospholipids that are changed under physiological pH due to interaction, absorption and distribution of polyphenols can be changed, too. To predict this interaction immobilized artificial membrane (IAM) columns are used (Figure 3). The measurement of these interactions as well as for RP-HPLC is  $\log k_{wIAM}$ .

To assess the binding of polyphenols to the plasma protein columns containing human serum albumin (HSA) and alfa1-glycoprotein acid are used. The percentage of polyphenol bounded to HSA is calculated using the equation:

$$\%HSA = 100 \frac{C_{stationary}}{C_{stationary} + C_{mobile}} = 100 \frac{k}{k+1}. \quad (8)$$

If the polyphenol shows a binding of over 95% it can potentially interact with other drugs of high affinity to HSA as the concentration of active (free) drug can increase significantly (e.g. 5% to 8% is increase of 60% in active form of drug).

### **Quantitative Structure-Activity Relationship [16]**

Quantitative Structure-Activity Relationship (QSAR) represents the analysis in which the structure of molecules is quantitatively correlated to biological activity. QSAR is now an irreplaceable technique in the rational design of drugs. The main postulate of QSAR is: biological activity is the consequence of chemical structure. To characterize the structure, different topological indices and three-dimensional (3D) descriptors are used. The most common topological indices are described in Table 1.

Besides QSAR, evolution in rational design has been achieved by using software for assessing physicochemical properties and parameters of bioavailability.

## **Review**

### **Thin Layer Chromatography**

Thin layer chromatography of standard solutions of flavonoids and phenolic acids was performed on silicagel plates 60 F<sub>254</sub> using different mobile phases (Table 2) and was visualized under short and long ultraviolet (UV) light after spraying with 1% AlCl<sub>3</sub>. The optimization was done using program KT1 [17]. Eleven mobile phases, for separation of flavonoids and phenolic acids were analyzed. The most suitable mobile

phases were number 7 (*n*-hexane : ethyl acetate : acetic acid (glac.) = 31 : 14 : 5 v/v) and number 11 (chloroform : methanol : formic acid = 44 : 3.5 : 2.5 v/v), as shown by dendrogram (Figure 4) and confirmed by values of discriminating power and the average information content (Table 3) [17]. Using mobile phases 11 and 7 we succeeded to identify up to 9 polyphenols in samples of Croatian propolis.

Analysis of flavonoids and phenolic acids in propolis was performed by two-dimensional thin layer chromatography (2D-TLC) using optimized chromatographic systems and detection and quantification was done using CAMAG Reprostar 3 densitometer with CAMAG VideoScan TLC/HPTLC evaluation software. The absorbance was recorded at 254 and 366 nm, the later being the wavelength of quantification [18]. Since propolis is a rather complex mixture for simple TLC, 2D-TLC analysis of propolis samples previously optimized chromatographic systems were used. Plate and densitograms of standard solution of caffeic acid and propolis sample from Peruća are presented in Figure 5.

2D-TLC was shown to be a suitable method for quantitative determination of flavonoids and phenolic acids in Croatian propolis samples. The linear relationship between absorbance and mass of standard was established:

$$A = 9706.6 + 4487.0 m; r = 0.98 \quad (9)$$

The presence of coumaric and cinnamic acids could not be established, as these compounds do not form fluorescent complexes with  $\text{AlCl}_3$ .

For the quantification purpose high performance thin layer chromatography (HPTLC) under previously optimized conditions was used. Seven polyphenols (Figure 6) have been quantified and the method was validated for a routine control of propolis tinctures [19]. HPTLC method was used for quantification and the results of

validation are shown in Table 4. HPTLC method was applied to 7 propolis samples and results are shown in Table 5.

For the investigation of ADME for flavonoids reversed-phase thin layer chromatography was used. Chromatography was performed on RP-18F<sub>254s</sub> plates with binary mobile phase prepared from methanol and phosphate buffer. The pH was at least 2 units lower than p*K*<sub>a</sub> value of phenolics, and organic modifier was varied from 30 to 80% in 5% increments. ADME data was calculated using the following programs: CLOGP, KowWin, Molinspiration, MolSoft, XLOGP, Osiris, VEGA ZZ, DRAGON, ChemSilico, ALOGPS, Interactive Analysis, SPARC [20]. Regular retention behaviour was observed for all the polyphenols investigated – a liner decrease of retention with increased concentration of organic modifier of mobile phase, with standard deviation of less than 0.1. To compare values of retention of polyphenols *R*<sub>Mw</sub> values were extrapolated to 100% buffer in mobile phase by using the equation:

$$R_M = R_{Mw} + b\phi, \quad (10)$$

where  $\Phi$  is the concentration of organic modifier. *R*<sub>Mw</sub> values of flavonoids were in the range 2.309-3.580 and phenolic acids 1.318-2.176. The correlation between *R*<sub>Mw</sub> and  $\Phi_0$  (*R*<sub>M</sub> = 0) was established:

$$R_{Mw} = 4.1854(\pm 0.2238) - 0.0569(\pm 0.0079)\phi_0, \quad (11)$$

$$n = 30, r = 0.941, s = 0.226, F = 216.$$

It is evident that *R*<sub>Mw</sub> and  $\Phi_0$  are well correlated but are not identical. This is the case with isomers sakuranetin and isosakuranetin that have the same chromatographic lipophilicity *R*<sub>Mw</sub> = 3.111, but different  $\Phi_0$ , 19.61 and 20.22 respectively.

Most of the used programs are “blind” to this difference occurring between structural isomers, and for MLOGP even different classes of flavonoids could not be distinguished: galangin (flavonol) and apigenin (flavone).

Besides lipophilicity, that is the main determinant for ADME, using ChemSilico software plasma protein binding, blood-brain barrier penetration and human absorption were predicted (Table 6).

BBB penetration was positive only for flavone and flavanone, as these do not have hydroxyl groups. Human absorption was the lowest for myricetin (58%).

Surprisingly, correlation of  $R_{Mw}$  and calculated log  $P$  values was rather poor, and the best correlations were found between  $\Phi_0$  and XLOGP values for flavonols, and human intestinal absorption calculated by ChemSilico and chromatographic  $R_{Mw}$  (for flavanones) and  $\Phi_0$  (for phenolic acids) values.

Based on the procedure developed by Ramić *et al.* [21] chemical interactions were studied using the optimized chromatographic system for polyphenols. Interactions of 19 polyphenols and over-the-counter (OTC) analgesic (acetylsalicylic acid, paracetamol, diclofenac, ibuprofen and ketoprofen) and vitamins (C and E) were examined. Interactions were detected between acetylsalicylic acid and acacetin, galangin, chrysin, pinocembrin and cinnamic acid; ibuprofen and flavone, ketoprofen and pinocembrin, vitamin C and myricetin, morin and quercetin; and vitamin E with flavanone. Example of chromatogram obtained is shown on Figure 7.

These chemical interactions cannot be replaced *in vivo* analysis, but rather be used as basis for further research of phenolics in biological systems.

Band-blot test was used to evaluate antioxidant activity of propolis samples compared to standard mixture of caffeic acid, galangin and pinocembrin using stable free radicals: 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS<sup>+</sup>) and the

2,2-diphenyl-1-picryl-hydrazyl (DPPH') [22]. Band-blot test was performed instead of original dot-blot test using the stable free radical of DPPH'. The reason for replacing dots with bands is equal to the distribution of samples in a line instead of over-concentrated centre of spot that can not be easily quantified. Antioxidant efficacy (AOE) was expressed as the slope of the dose-response curve of decrease in DPPH' radical absorbance was used. The chromatogram for standard mixture is shown on Figure 8. Band-blot test has proven a rather quick method for assessing antioxidant activity of a great number propolis samples as the only application of samples and visualization after spraying was done, taking less than 10 minutes.

For wine samples the same procedure of optimization was used to select the most appropriate mobile phase (benzene : ethyl acetate : formic acid = 30 : 15 : 5 v/v). Six phenolic compounds were identified in the tested samples. The plates are shown on Figure 9 [23]. TLC quantification of polyphenols in wine was performed using CAMAG system. The substances were identified on basis of  $R_F$  values and UV spectra. Densitometric quantitative analysis of gallic acid in wine extract is presented on Figure 10, as an example.

### **High Performance Liquid Chromatography**

HPLC analysis of propolis was conducted on Agilent 1100 using X-bridge C18 column. Binary mobile phase of eluents A (10 mM ammonium-formiate in water) and B (10 mM ammonium-formiate in methanol) both of pH = 7 (adjusted with formic acid) was used with the gradient stated in Table 7 [22]. Flow rate was 0.5 mL/min at room temperature. 21 standards of polyphenols were used; their identification was based on retention time and UV spectrum, and quantification was done at 270 nm.

Reverse phase HPLC was used for the lipophilicity/hydrophobicity of flavonoids and phenolic acids. Mobile phase A was 2% acetic acid in water and B 2% acetic acid in

methanol. Gradient method started with 95% of mobile phase A and 5% of mobile phase B that was linearly increased to 100% during 20/60 minutes [24]. As shown on chromatogram, phenolic acids as more polar substances were first eluted from the column, followed by more hydrophilic and lipophilic flavonoids. HPLC results were used to identify flavonoids and phenolic acids present in propolis samples, to quantify the content of polyphenols, to explain the antioxidant activity *in vitro* and *in vivo* as well as they were used for calculations of pharmacokinetic parameters [22].

For the IAM-HPLC analysis IAM.PC.DD 2 column was used. Mobile phase was phosphate buffer modified with different volumes of methanol. For the HSA-HPLC analysis Chiral HSA column and gradient of phosphate and 2-propanol were used. IAM-HPLC analysis showed that ionization of polyphenolic under physiological pH was influencing the retention of the substance due to ion-ion interaction. The strongest interactions were in the group of flavonoids (kaempferide  $\log k_{wIAM} = 3.581$ ) and lower for phenolic acids ( $\log k_{wIAM} = 1.35-1.68$ ).

From the results of HSA-HPLC analysis it has to be noticed that all the flavonoids have percentage of bonded fraction greater than 95%, which can have repercussions to the interactions with drugs. Binding to HSA was higher for myricetin (%HSA = 99.8) from the group of flavonoids and the lowest for caffeic acid (%HSA = 81.8) from the phenolic acids analyzed, both having the greatest number of hydroxyl groups. Enantiomers could have been differentiated using this method, as expected. This is the case with flavanons shown on Figure 11.

HPLC analysis of wine was conducted on Agilent 1100 using Zorbax StableBond C18 column [5]. Binary mobile phase of eluents A (phosphoric acid (c = 0.02 M, pH = 3.0 adjusted with triethylamin) : methanol = 90 : 10 v/v) and B (phosphoric acid (c = 0.02 M, pH = 3.0 adjusted with triethylamin) : methanol = 10 : 90 v/v) was used with the

gradient stated in Table 8. Flow rate was 1 mL/min at 35 °C. 15 standards of polyphenols were used; their identification was based on retention time and UV spectrum, and quantification was done at 280 nm. HPLC analysis of samples of wine has shown discrepancy in results compared to TLC. This has been described in literature [25,26] and can be explained by lower limit of detection and possibility of overlapping the peaks in HPLC. To illustrate this comparative results for Pelješac sample are shown in Table 9. Chromatogram of wine samples from Croatia obtained by HPLC analysis is shown on Figure 12.

### **Spectrophotometry**

The Folin-Ciocalteu method was used for quantification of total polyphenols according to the method described by Slinkard and Singleton [27]. The method is based on the reduction of  $\text{MoO}^{4+}$  to  $\text{MoO}^{3+}$  that is detected by color change from yellow to blue; measured at 765 nm. The results were expressed as equivalents of gallic acid from the calibration curve.

The results for determination of total polyphenols, according to the Folin-Ciocalteu method have shown that white wines (Malvazija (*malv*), Traminac (*tram*), Pelješac (*pelj*)) have a lower content of polyphenols [5]. Red wine contains more polyphenols than white wine. This may be ascribing to the fact that making of white wine requires the removal of the skins after the grapes are crushed. Wines containing higher contents of polyphenols are those from the south of Croatia (Plavac (*plav*), Postup (*post*), Dingač (*ding*)) (Figure 13). These findings have been confirmed by TLC analysis.

Free stable radicals DPPH<sup>•</sup> and ABTS<sup>•+</sup> were used to assess the antioxidant activity. The method is based on lost of violet/blue color of radicals in presence of substance capable to accept free electron. Discoloration was measured on spectrophotometer at 518 and 730 nm respectively [22]. The spectrophotometry assays, using stable free

radicals to assess the antioxidant activities have proven reliable and precise. As most of the suggested procedures could not differentiate similar samples, kinetics of different concentrations of propolis samples has been studied and antioxidant efficiency was introduced (AOE). DPPH<sup>•</sup> radical was more stable than ABTS<sup>•+</sup> and results of the DPPH<sup>•</sup> assay showed better reproducibility. Based on the content of flavonoids and phenolic acids identified and quantified by HPLC analysis we obtained the following correlation:

$$AOE_{DPPH} = 0.5637 + 0.0285c_{phenolic\ acids} + 0.0063c_{flavonoids}, \quad (12)$$

$$n = 5, r = 0.878, s = 0.4713, F = 3.3623$$

### **Quantitative Structure-Activity Relationship**

Experimental data used for QSAR analysis was taken from the literature. Wiener index, connectivity index, Balaban index, Balaban-type indices from atomic number, mass, van der Waals, electronegativity and polarizability weighted distance matrix, information-theoretic index, Shultz index, together with molecular weight, n-octanol/water partition coefficient, van der Waals volume, molar refractivity and polar surface area of polyphenols were calculated using TAM [28], HyperChem 8.0 Evaluation software, PCLIENT, Dragon 3.0. Four groups of 3D descriptors were used: geometrical, GETAWAY (Geometry, Topology and Atom Weights Assembly), 3D-MORSE and RDF (Radial Distribution Function). Linear, polynomial and multiple linear regression analysis was conducted using Statistica 6.0 [16].

Statistically significant QSAR models for lipid peroxidation inhibiting effects of flavonoids were obtained by polynomial and multiple regression using lipophilicity, Balaban index, Balaban-type index and 3D GETAWAY descriptor. The reason for selecting the 3D descriptors was the possibility of discriminating stereoisomers, like catechin and epicatechin (Figure 14).

The best model that includes 3D descriptor  $H_7(p)$  in combination with Balaban-type index from mass-weighted distance matrix ( $J_m$ ) was chosen by the best-subset regression:

$$IC_{50} = 320.3(\pm 247.9) + 17.46(\pm 1.92)J_m + 19.63(\pm 2.59)H_7(p), \quad (13)$$

$$n = 8, r^2 = 0.95, S = 7.87, F = 47.66.$$

$H_7(p)$  is an autocorrelation descriptor calculated for 3D-spatial molecular geometry based on lag (topological distance) and weighted by atomic polarizabilities. This descriptor belongs to the group of H-GETAWAY descriptors that have been calculated from the molecular influence matrix H. These descriptors are sensitive to significant conformational changes and to the bond lengths that account for atom types and bond multiplicity.

This QSAR investigation of wine polyphenols could be extended to a larger number of substances, including anthocyanins and dimeric procyanidins as the most common polyphenols in wine.

### **Clinical Trial [29]**

An *in vivo* study has been conducted on 47 healthy women and men in order to investigate whether daily intake of powdered propolis extract during 30 days has any influence on antioxidative status based on the following blood parameters: activity of superoxide dismutase, glutathione peroxidase and catalase, concentration of plasma malondialdehyde, total cholesterol, low- and high-density lipoprotein cholesterol, triglycerides, glucose, uric acid, ferritin and transferrin, together with routine red blood cell parameters.

This was the first reporting of the effects of prolonged propolis supplementation on redox-status of human organism. The benefit of propolis use was only shown in male population demonstrating reduction in free-radical-induced lipid peroxidation as well

as increase in activity of superoxide dismutase. A 23.2% decrease in malonaldehyde (degradation product of peroxidation of polyunsaturated fatty acids) concentration and a 20.9% increase in superoxide dismutase activity (first and most important line of antioxidant enzyme defense) were observed (Figure 15).

Absence of the same results in women was probably due to uncoordinated menstrual cycles. Hence, estrogens are powerful antioxidants, modulators of antioxidant enzyme expression and levels of lipid peroxidase and lipoproteins.

## **Summary and conclusion**

In the last decade interest on nutraceuticals and natural medicinal products is constantly growing. The market is full of antioxidant formulations of wide variety of sources. For the registration of nutraceutical as natural medicinal product analytical procedures have to be developed, product has to be standardized and their functionality and beneficial effects have to be demonstrated. The goal of our studies was standardization of Croatian propolis extracts as a rich sample of flavonoids and phenolic - active components to which pharmacological (antioxidant) activity is attributed. Parallel research on wine samples was done, as wine represents another sample rich in polyphenolics that has been used as constituents e.g. iron wine, tonics.

### **Identification and quantification of flavonoids and phenolic acids**

Although the methods for identification and quantification of propolis samples using HPLC combined with mass spectrometry (MS) and nuclear magnetic resonance (NMR) have been developed and alternative was provided in capillary electrophoresis coupled with MS, these are rather expensive, not readily available and inappropriate for routine analysis [30,31]. Thus we have decided for TLC method combined with HPLC-diode array detection (DAD) method. Out of 11 mobile phases available from

literature based on numerical taxonomy chloroform : methanol : formic acid in volume ratio 44 : 3.5 : 2.5 was selected as the most appropriate for TLC identification of 9 polyphenols [17]. Better separation for identification was achieved using 2D-TLC chromatography, although for quantification we decided for high performance TLC plates using which clear identification and quantification of 7 polyphenols (namely: *p*-coumaric acid, caffeic acid, chrysin, tectochrysin, pinocembrine-7-methyl ether, isoferulic acid) was performed [18,19]. Major limiting factor of TLC is length of the plate, so for better separation HPLC was used.

Total content of polyphenols was determined using Folin-Ciocalteu method. As we have done identification using liquid chromatography we did not separately analyze total content of phenolic acids and flavonoids which can be done using well known procedures described in Ph. Eur. and Christ and Müller, respectively [32,33].

#### **Pharmacokinetic (ADME) parameters of flavonoids and phenolic acids**

Passive transport through membranes is mainly determined by the lipophilicity of the substance. The most commonly used experimental value of lipophilicity is chromatographic parameter  $R_M$ . This parameter was determined by TLC as well by HPLC and to assess lipophilicity within the group of polyphenolics. As this parameter cannot distinguish isomers, additional parameter  $\Phi_0$  was used differentiating sakuranetin and isosakuranetin.

Using ChemSilico software plasma protein binding of flavonoids was found rather high (85-96%) compared to phenolic acids (70-86%). High values of flavonoid binding are in accordance to the experimental data e.g. for quercetin (99%) [34]. Passage through blood-brain barrier was negative which is expected for compounds having acidic phenolic and carboxylic groups. Absorption was mainly greater than

60%, but this has to be taken with dose of doubt, as experimental data is contradictory [35].

Chemical interaction with most commonly used non-steroidal anti-inflammatory drugs (NSAID) and vitamins were characterized on TLC plate. The interactions were most common with acetylsalicylic acid. At the moment this has no practical value as further *in vivo* studies have to demonstrate applicability of this model [21].

Interactions of polyphenolics with artificial membranes were probably based on ion-ion and passive diffusion as interaction showed greater partition coefficient for flavonoids than phenolic acids. Binding on HSA-column was higher compared to ChemSilico predictions and more in accordance to experimental data obtained *in vivo*.

#### **Antioxidant activity of flavonoids and phenolic acids**

Antioxidant activity of polyphenolics is well known. To assay the antioxidant activity different methods are used: electron paramagnetic resonance (EPR) spectroscopy, chemiluminescence, enzyme assays... Most often used and readily available is spectrophotometric method using stable free radicals. DPPH<sup>·</sup> and ABTS<sup>·+</sup> were used for spectrophotometry as well TLC *in situ* for analysis of total antioxidant capacity of propolis. The most active propolis comes from coastal Croatia [22].

QSAR analysis of polyphenolics based on Trolox test data from literature pointed out that antioxidant activity of polyphenols as hydrogen donating free radical scavengers, is closely related to their chemical structure, especially with the number and arrangement of free hydroxyl groups of polyphenol skeleton [16].

*In vivo* study of propolis prolonged used showed beneficial in male population demonstrating reduction in free-radical-induced lipid peroxidation as well as increase in activity of superoxide dismutase. Production of malonaldehyde (degradation product of peroxidation of polyunsaturated fatty acids) reduced and activity of

superoxide dismutase (first and most important line of antioxidant enzyme defense) was increased [29].

Antioxidant supplements are flooding the market. Asking pharmacist for a new product - extract of the plant coming from exotic country, will usually end with the answer “its antioxidant and thus good for your health”. The necessity of standardization natural antioxidant products made us write this minireview-providing basis for standardization of natural antioxidant products rich in polyphenols using simple and readily available techniques based on our research on propolis and wine.

## **Abbreviations**

ABTS: 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid; ADME: absorption, distribution, metabolism, excretion; AOE: antioxidant efficacy; AU: arbitrary unit; BBB: blood brain barrier; 3D descriptors: three-dimensional descriptors; 2D-TLC: two-dimensional thin layer chromatography; DAD: diode array detection; *DP*: discriminating power; DPPH: 2,2-diphenyl-1-picryl-hydrazyl; EPR: electron paramagnetic resonance; GAE: gallic acid equivalent; GETAWAY: geometry, topology and atom weights assembly; GIT: gastrointestinal tract; HSA: human serum albumin; HPTLC: high performance thin layer chromatography; IAM: immobilized artificial membrane; MS: mass spectrometry; NMR: nuclear magnetic resonance; NSAID: non-steroidal anti-inflammatory drugs, OTC: over-the-counter; RDF: radial distribution function;  $R_F$ : retention factor;  $R_M$ : chromatographic parameter of lipophilicity; RP-HPLC: reverse phase high performance liquid chromatography; RP-TLC: reverse phase thin layer chromatography; SOD: superoxide-dismutase; UV: ultraviolet.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

All authors contributed equally to this work, read and approved the final manuscript.

## Acknowledgements

This work was supported by the Ministry of Science, Education and Sports of the Republic of Croatia (projects No. 006-0061117-1237 - MMŠ and 006-0061117-1239 - ŽM). The authors wish to thank all the associates involved in these as well as antecedent projects.

## References

1. Bradamante V, Lacković Z: *Oxidative Stress and Efficiency of Antioxidants*. Zagreb: Medicinska naklada; 2001.
2. Havsteen B: **Flavonoids, a class of natural products of high pharmacological potency**. *Biochem Pharmacol* 1983, **7**:1141-1148.
3. Smirnoff N: *Antioxidants and Reactive Oxygen Species in Plants*. Oxford: Blackwell Publishing; 2005.
4. Castalado S, Capasso F: **Propolis, an old remedy used in modern medicine**. *Fitoterapia* 2002, **73**(Suppl 1):S1-S6.
5. Rastija V, Medić-Šarić M: **Polyphenolic composition of Croatian wines with different geographical origins**. *Food Chem* 2009, **115**:54-60.
6. Kaštelan-Macan M, Medić-Šarić M, Turina S: *Planar Chromatography*. Zagreb: Faculty of Pharmacy and Biochemistry; 2006.

7. Medić-Šarić M, Maleš Ž, Stanić G, Šarić S: **Evaluation and selection of optimal solvent and solvent combinations in thin-layer chromatography of flavonoids and of phenolic acids of *Zizyphus jujuba* Mill.** *Croat Chem Acta* 1996, **69**:1265-1274.
8. Medić-Šarić M, Stanić G, Maleš Ž, Šarić S: **Application of numerical methods to thin-layer chromatographic investigation of the main components of chamomile (*Chamomilla recutita* (L.) Rauschert) essential oil.** *J Chromatogr A* 1997, **776**:355-360.
9. Medić-Šarić M, Maleš Ž: **Application of numerical methods to the TLC analysis of flavonoids and phenolic acids of *Rosmarini folium*.** *J Planar Chromatogr* 1997, **10**:182-187.
10. Maleš Ž, Medić-Šarić M, Bucar F: **Flavonoids of *Guiera senegalensis* J.F.GMEL. – thin-layer chromatography and numerical methods.** *Croat Chem Acta* 1998, **71**:69-79.
11. Medić-Šarić M, Maleš Ž, Šarić S, Debeljak Ž: **Use of information theory and numerical taxonomy methods for evaluating the quality of thin-layer chromatographic separations of flavonoids and phenolic acids of *Rhamni cathartici fructus*.** *J Liq Chrom & Rel Technol* 1999, **22**:83-103.
12. Medić-Šarić M, Maleš Ž: **Selection of an optimal set of solvents in thin-layer chromatography of flavonoids and phenolic acids of *Lavandulae flos*.** *Pharmazie* 1999, **54**:362-364.
13. Maleš Ž, Medić-Šarić M: **Investigation of the flavonoids and phenolic acids of *Sambuci flos* by thin-layer chromatography.** *J Planar Chromatogr* 1999, **12**:345-349.

14. Maleš Ž, Medić-Šarić M: **Optimization of flavonoids and phenolic acids of *Helleborus atrorubens* Waldst. et Kit.** *J Pharm Biomed Anal* 2001, **24**:353-359.
15. van de Waterbeemd H, Hans Lennernäs H, Per Artursson P: *Drug Bioavailability: Estimation of Solubility, Permeability, Absorption and Bioavailability*. Weinheim: Wiley-VCH; 2003.
16. Rastija V, Medić-Šarić M: **QSAR Study of antioxidant activity of wine polyphenols.** *Eur J Med Chem* 2009, **44**:400-408.
17. Jasprica I, Smolčić-Bubalo A, Mornar A, Medić-Šarić M: **Investigation of the flavonoids in Croatian propolis by thin-layer chromatography.** *J Planar Chromatogr* 2004, **17**:95-101.
18. Medić-Šarić M, Jasprica I, Mornar A, Smolčić-Bubalo A, Golja P: **Quantitative analysis of flavonoids and phenolic acids in propolis by two-dimensional thin layer chromatography.** *J Planar Chromatogr* 2004, **17**:459-463.
19. Cvek J, Medić-Šarić M, Jasprica I, Mornar A: **High-performance thin-layer chromatographic analysis of the phenolic acid and flavonoid content of Croatian propolis samples.** *J Planar Chromatogr* 2007, **20**:429-435.
20. Mornar A, Medić-Šarić M, Jasprica I: **ADME data for polyphenols characterized by reversed-phase thin-layer chromatography.** *J Planar Chromatogr* 2006, **19**:409-417.
21. Ramić A, Medić-Šarić M, Turina S, Jasprica I: **TLC detection of chemical interaction of vitamins A and D with drugs.** *J Planar Chromatogr* 2006, **19**:27-31.

22. Jasprica I, Bojić M, Mornar A, Bešić E, Bućan K, Medić-Šarić M: **Evaluation of antioxidant activity of Croatian propolis samples using DPPH<sup>•</sup> and ABTS<sup>•+</sup> stable free radical assays.** *Molecules* 2007, **12**:1006-1021.
23. Rastija V, Mornar A, Jasprica I, Srećnik G, Medić-Šarić M: **Analysis of phenolic components in croatian red wine by thin-layer chromatography.** *J Planar Chromatogr* 2004, **17**:26-31.
24. Medić-Šarić M, Mornar A, Jasprica I: **Lipophilicity of flavonoids and phenolic acids determined by shake-flask and high performance liquid chromatography methods.** In *Proceedings of the 6th AFMC International Medicinal Chemistry Symposium: 8-11 July 2007; Istambul.* Barcelona: Prous science; 2007:133-133.
25. Shi R, Schwedt G: **HPLC and TLC analysis of phenolic compounds in wine.** *Deut Lebensm-Rundsch* **1** 1995: 14-17.
26. de Rijke E, Out P, Niessen WMA, Ariese F, Gooijer C, Brinkman UATh: **Analytical separation and detection methods for flavonoids.** *J Chromatogr A* 2006, **1112**:31-36.
27. Slinkard K, Singleton VL: **Total phenol analysis: automation and comparison with manual methods.** *Am J Enol Vitic* 1977, **28**:49-55.
28. Vedrina M, Marković S, Medić-Šarić M, Trinajstić N: **TAM: a program for the calculation of topological indices in QSPR and QSAR studies.** *Computers Chem* 1997, **21**:355-361.
29. Jasprica I, Mornar A, Debeljak Ž, Smolčić-Bubalo A, Medić-Šarić M, Mayer Lj, Romić Ž, Bućan K, Balog T, Sobočanec S, Šverko V: **In vivo study of propolis supplementation effects on antioxidative status and red blood cells.** *J Ethnopharmacol* 2007, **110**:548-554.

30. Cuesta-Rubio O, Piccinelli AL, Fernandez MC, Hernandez IM, Rosado A, Rastrelli L: **Chemical characterization of Cuban propolis by HPLC-PDA, HPLC-MS, and NMR: the brown, red, and yellow Cuban varieties of propolis.** *J Agric Food Chem* 2007, **55**:7502-7509.
31. Gomez-Romero M, Arraez-Roman D, Moreno-Torres R, Garcia-Salas P, Segura-Carretero A, Fernandez-Gutierrez A: **Antioxidant compounds of propolis determined by capillary electrophoresis-mass spectrometry.** *J Sep Sci* 2007, **30**:595-603.
32. Council of Europe: *European Pharmacopoeia*. Strasbourg; 2007.
33. Christ B, Müller KH: **Zur serienmäßigen bestimmung des gehaltes an flavonol derivaten in drogen.** *Arch Pharm* 1960, **293**:1033-1042.
34. Hollman PC, Katan MB: **Absorption, metabolism and health effects of dietary flavonoids in man.** *Biomed Pharmacother* 1997, **51**:305-310.
35. Walle T: **Absorption and metabolism of flavonoids.** *Free Radic Biol Med* 2004, **36**:829-837.

## Figures

**Figure 1 - Basic structures of flavonoids (A) and phenolic acids (B-cinnamic acid, C-benzoic acid).**

**Figure 2 - The map of Croatia. Areas of origin of propolis samples (marked by numbers).**

**Figure 3 - The structure of stationary phase of immobilized artificial membrane – phosphatidylcholine – drug-discovery column (IAM.PC.DD).**

**Figure 4 - Dendrogram for eleven TLC mobile phases used for determination of polyphenols in samples of Croatian propolis.**

**Figure 5 - 2D-TLC of caffeic acid in propolis sample from Peruča and standards of caffeic acid.**

The plate was first developed in the direction 1 then 2 using mobile phase 1 (n-hexane : ethyl acetate : glacial acetic acid = 31 : 14 : 5 v/v) and mobile phase 2 (chloroform : methanol : formic acid = 44 : 3.5 : 2.5 v/v). Bands were visualized under short and long wave UV light and under long wave UV light after spraying with 1%  $\text{AlCl}_3$  ethanol solution. Concentration obtained from chromatograms:  $a = m_{\max} = 2.50 \mu\text{g}$ ,  $b = m_{\min} = 0.83 \mu\text{g}$ ,  $c = m_{\text{mid}} = 1.67 \mu\text{g}$ ,  $\text{PS} = m_{\text{propolis sample}} = 2.91 \mu\text{g}$  [18].

**Figure 6 - Structures of flavonoids and phenolic acids quantified in propolis tinctures.**

**Figure 7 - Chromatogram of interaction of flavonoid pinocembrin and ketoprofen.**

Interaction of flavonoid pinocembrin and ketoprofen obtained under long wave UV light. SF - standard of flavonoid (0.33 mg/mL), A, B, C - interaction lines of pinocembrin and ketoprofen in concentration 0.5, 1.0 and 1.5 mg/mL respectively, SL - standard of ketoprofen in the same concentration as in A, B and C.

**Figure 8 - Band-blot test of polyphenols present in propolis using the stable free radical DPPH.**

Chromatogram obtained after spraying the layer (applied polyphenol standard mixture bands in the range from 45 (track 1) to 10  $\mu\text{g/mL}$  (track 8)) with 0.3 mM solution of DPPH $\cdot$  radical. *AOE* of standard mixture is 1.89 ( $H = 1.89c + 22.856$ ,  $r^2=0.974$ ) [22].

**Figure 9 - Chromatograms of extracts of red wines “Merlot” (1) and “Frankovka” (2) by use of mobile phase benzene : ethyl acetate : formic acid = 30 + 15 + 5 v/v/v.**

a) recorded under UV light  $\lambda = 254$  nm; b) recorded under UV light  $\lambda = 366$  nm, after spraying with 1% ethanol solution of  $\text{AlCl}_3$ . The standard substances used: A – gallic acid, B – caffeic acid, C – apigenin, D – kaempferol, E – *p*-coumaric acid, F – naringenin [23].

**Figure 10 - TLC identification and quantification of gallic acid.**

Photograph of the chromatographic plate recorded at UV light  $\lambda = 254$  nm, for the purpose of quantification of gallic acid in the sample of wine, with the spectrum of marked peaks obtained using CAMAG densitometer.

**Figure 11 - Structure and differences in HSA binding of enantiomers.**

**Figure 12 - HPLC chromatogram of wine sample Postup registered at 280 nm.**

**Figure 13 - Total polyphenols in wine samples as equivalents of gallic acid (GAE, mg/L) [5].**

Codes for wine samples: *malv* - Malvazija; *tram* – Traminac; *pelj* – Pelješac; *zwei* – Zweigelt; *merl* – Merlot; *fran* – Frankovka; *cabs* – Cabernet Sauvignon; *borg* – Borgonja; *pinc* – Pinot crni; *ding* – Dingač; *post* – Postup; *plav* – Plavac.

**Figure 14 - Structures of catechin and epicatechin.**

**Figure 15 - Antioxidant activity of propolis *in vivo*.**

Activity of superoxide-dismutase (SOD) (U/g Hb) during the study: a) in the men test group and b) in the women test group. Data are expressed as means  $\pm$ S.D.; B (1) represents the baseline on the 15<sup>th</sup> day prior to the study and PS (30) represents

propolis supplementation on the 30<sup>th</sup> day. The symbol (\*) denotes statistically significant change [29].

## Tables

**Table 1 - The most commonly used topological indices.**

Index	Expression for calculation	Observation
Wiener index	$W(G) = \frac{1}{2} \sum_{i,j} D_{i,j}$	$D_{i,j}$ represents off-diagonal elements of matrix which stands for the shortest distance in term on number of bonds between atom $i$ and $j$
Valence connectivity index	$\chi^v(G) = \sum_{i,j} \frac{1}{\sqrt{v_i v_j}}$	$v_i$ and $v_j$ are weights (valence delta values) of vertices $i$ and $j$ making up edge in vertex weighted graph $G$
Balaban index	$J(G) = \frac{E}{\mu + 1} \sum_{edges} \frac{1}{\sqrt{ds_i ds_j}}$	the average distance sum connectivity; where $E$ is the number of edges, $\mu$ is cyclomatic number of $G$ and $ds_i$ is a distance sum
Information-theoretic index	$I(G) = - \sum_i \frac{2N_i}{N(N-1)} \log_2 \frac{2N_i}{N(N-1)}$	modified Shannon's equation where $n$ is the number of different sets of elements, $N_i$ is the number of elements in the $i$ -th set of elements and the sum is over all sets of elements
Shultz index	$MTI = \sum_i e_i$	molecular topological index (MTI) is based on adjacency matrix ( $A$ ), the distance matrix ( $D$ ) and the valency matrix ( $v$ ); the of elements $e_i$ of the row matrix $v[A+D]$ gives Shultz index

**Table 2 - The mobile phases studied.**

No	Solvents	Proportions (v/v)
1	toluene - ethyl acetate - formic acid (98-100%)	36 : 12 : 5
2	cyclohexane - ethyl acetate - formic acid (98-100%)	30 : 15 : 5
3	toluene - ethyl acetate - glacial acetic acid	36 : 12 : 5
4	cyclohexane - ethyl acetate - glacial acetic acid	31 : 14 : 5
5	<i>n</i> -hexane - ethyl acetate - formic acid (98-100%)	31 : 14 : 5
6	toluene - acetone - formic acid (98-100%)	38 : 10 : 5
7	<i>n</i> -hexane- ethyl acetate - glacial acetic acid	31 : 14 : 5
8	petroleum ether (40-70 °C) - ethyl acetate - formic acid (98-100%)	30 : 15 : 5
9	carbon tetrachloride - acetone - formic acid (98-100%)	35 : 10 : 5

<b>10</b>	<i>n</i> -hexane - ethyl acetate - glacial acetic acid	30 : 20 : 1.5
<b>11</b>	chloroform - methanol - formic acid (98-100%)	44 : 3.5 : 2.5

**Table 3 - DP and I output data for error factor E = 0.03 for each mobile phase.**

Mobile phase	DP	I (bit)
1	0.8538	3.221
2	0.7836	2.735
3	0.8655	3.076
4	0.8947	3.616
5	0.8187	2.860
6	0.8480	3.011
<b>7</b>	<b>0.9298</b>	<b>3.682</b>
8	0.7895	3.050
9	0.9240	3.511
10	0.9181	3.471
<b>11</b>	<b>0.9415</b>	<b>3.827</b>

**Table 4 - Results of validation of HPTLC method [19].**

Validation data	Value
linearity	r > 0.99
precision	
repeatability of the sample application and intra-day precision	RSD < 3.8%
repeatability of peak-area measurement	RSD < 2.4%
accuracy	RSD < 6.5%
limit of detection	PME 7.5 ng/band IFA 7.5 ng/band CA 60 ng/band
limit of quantification	PME 22.5 ng/band IFA 22.5 ng/band CA 180 ng/band
robustness	
inter-day precision and stability of standard solutions	RSD < 5% (two days)
stability on the plate	RSD < 3.4% (up to 2 hours)*

effect of temperature

RSD < 4.6% (20/26 °C)

selectivity

absorption spectra matching > 0.99

Legend:

\*only caffeic acid degraded up to 9.4% in 120 minutes, but as the analysis lasts less, this was not of great significance; CA: caffeic acid; IFA: isoferulic acid; PME: pinocembrin-7-methylether; RSD: relative standard deviation.

**Table 5 - Phenolic acid and flavonoid content (expressed as mean of mass concentration in mg/mL) of the Croatian propolis samples analyzed [19].**

propolis sample	polyphenolic				pinocembrin		isoferulic acid
	<i>p</i> -coumaric acid	caffeic acid	chrysin	tectochrysin	pinocembrin	-7-methyl ether	
Veliki Zdenci	0.3223+1.32	0.1204+0.99	-	-	-	-	-
Sisak	0.1331+8.22	0.2675+0.22	0.1423+4.09	0.0845+2.90	-	-	-
Labin	0.1464+3.82	0.3000+0.74	0.3987+3.30	-	-	-	-
Čisla (Omiš)	0.2047+2.09	0.4123+0.51	0.1247+1.84	-	-	0.0593+7.36	-
Pelješac	-	0.4196+0.38	0.4896+1.46	0.6775+1.45	0.8454+2.13	0.7452+0.84	0.5068+0.59
Metković	0.2968+1.04	0.3538+0.79	0.5616+2.62	1.1816+0.81	0.6725+1.54	1.1621+0.85	0.4376+2.42
Mixture of propolis	0.1908+2.14	0.3398+0.92	0.6237+4.47	0.4179+1.71	1.1465+3.24	0.2305+5.49	0.4109+4.54

**Table 6 - Results obtained using ChemSilico software [20].**

ADME data	Results
plasma protein binding	
- flavonoids	85.41% (myricetin) – 95.65% (flavone)
- phenolic acids	70.22% (sinapic acid) – 85.99% (cinnamic acid)
blood-brain barrier penetration	negative except flavon and flavanone
human absorption	significant

**Table 7 - Gradient of mobile phases used for HPLC analysis of propolis.**

time (min)	0	15	25	30	40	50	60	75	80
mobile phase									
%A	90	50	50	40	40	10	10	90	90
%B	10	50	50	60	60	90	90	10	10


**Table 8 - Gradient of mobile phases used for HPLC analysis of wine.**

time (min)	0	25	30
mobile phase			
%A	85	15	5
%B	15	85	95


**Table 9 - Comparison of the results obtained by HPLC and TLC analysis for Pelješac sample.**

	gentisic acid	gallic acid	cinnamic acid	<i>o</i> -coumaric acid	<i>m</i> -coumaric acid	<i>p</i> -coumaric acid	caffeic acid	ferulic acid	chlorogenic acid	catechin	quercetin	kaempferol	myricetin	quercitrin	apigenin	naringenin	<i>trans</i> -resveratrol
<b>HPLC</b>	na	8.4	na	-	-	-	5.0	0.7	-	4.1	4.1	0.4	-	-	-	-	0.8
<b>TLC</b>	48.23	8.92	36.56	na	-	-	3.50	0.93	na	na	5.06	-	na	na	-	-	na


Results expressed as mean of mass concentration in mg/L (na – not analyzed, - – not identified).



A



B



C

Figure 1





Figure 2








Figure 5




Figure 6



Figure 7

All trades @ 518 nm




Figure 8





Figure 9








Figure 10



pinocembrin  
( $\Delta\%HSA=4.7$ )



naringenin  
( $\Delta\%HSA=0.5$ )



hesperetin  
( $\Delta\%HSA=0.6$ )




Figure 12




Figure 13





Figure 14 catechin



epicatechin





Figure 15

**Additional files provided with this submission:**

Additional file 1: answer to reviewers.doc, 28K

<http://www.nutritionj.com/imedia/8348490932892653/supp1.doc>